

(80) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, DM, DO, DZ, EE, ES, FI, GB, GR, GD, GP, HK, HU, IL, IN, IS, JP, KG, KR, KZ, LK, LR, LU, LV, LY, MA, MD, MG, MK, MN, MX, NO, NZ, PL, PT, RO, RU, SE, SG, SI, SK, SL, SM, TN, TT, TZ, UA, UG, UZ, VU, YU, ZA, ZW, patent (TH, GM, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), patent (TH, GM, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BI, CF, CG, CI, CM, CA, GN, GW, ML, MR, NE, NG, TD, TO).

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LK	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	CA	Canada	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GR	Greece	MD	Republic of Moldova	TG	Togo
BB	Barbados	GI	Gibraltar	MG	Madagascar	TL	Timor-Leste
BD	Bangladesh	CN	China	MX	Mexico	TM	Turkmenistan
BE	Belgium	HR	Croatia	MY	Malaysia	TR	Turkey
BF	Burkina Faso	GR	Greece	ML	Mali	TT	Trinidad and Tobago
BG	Bulgaria	HU	Hungary	MR	Mauritania	UA	Ukraine
BJ	Benin	IE	Ireland	MN	Mongolia	UG	Uganda
BR	Brazil	IL	Israel	MT	Malta	US	United States of America
BY	Belarus	IS	Iceland	MY	Maldives	UZ	Uzbekistan
CA	Canada	IT	Italy	MX	Mexico	VN	Viet Nam
CF	Central African Republic	JP	Japan	NE	Niger	YU	Yugoslavia
CG	Congo	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CH	Switzerland	KE	Kenya	NL	Netherlands		
CI	Cote d'Ivoire	NP	Nepal	NZ	New Zealand		
CM	Cameroon	PR	Puerto Rico	PL	Poland		
CN	China	RK	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LJ	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

PEGYLATED INTERFERON ALFA- CCR5 ANTAGONIST COMBINATION HIV THERAPY

BACKGROUND OF THE INVENTION

The present invention relates to the use of a pegylated interferon-alfa and a CCR5 antagonist for the preparation of a medicament for the treatment of HIV-1 infections as well as HIV-1 and HCV co-infections in patients. The treatment involves administering a therapeutically effective amount of pegylated interferon-alfa in association with a therapeutically effective amount of a CCR5 antagonist sufficient to lower HIV-1-RNA.

The global health crisis caused by Human Immunodeficiency Virus-1 ("HIV-1"), the causative agent of Acquired Immunodeficiency Syndrome (AIDS), is unquestioned, and while recent advances in drug therapies have been successful in slowing the progression of AIDS, there is still a need to find a safer, more efficient, less expensive way to control the virus.

It has been reported that the CCR5 gene plays a role in resistance to HIV infection. HIV infection begins by attachment of the virus to a target cell membrane through interaction with the cellular receptor CD4 and a secondary chemokine co-receptor molecule, and proceeds by replication and dissemination of infected cells through the blood and other tissue. There are various chemokine receptors, but for macrophage-tropic HIV, believed to be the key pathogenic strain that replicates *in vivo* in the early stages of infection, the principal chemokine receptor required for the entry of HIV into the cell is CCR5. Therefore, interfering with the interaction between the viral receptor CCR5 and HIV can block HIV entry into the cell. The present invention relates to the use of small molecules which are CCR5 antagonists in association with pegylated Interferon-alfa to treat patients having HIV-1 infections.


A-M. Vandamme et al., Antiviral Chemistry & Chemotherapy, 9:187-203 (1998) disclose current clinical treatments of HIV-1 infections in man including at least triple drug combinations or so-called Highly Active Antiretroviral Therapy ("HAART"). HAART involves various combinations of nucleoside reverse transcriptase inhibitors ("NRTI"), non-nucleoside reverse transcriptase inhibitors

("NRTI") and HIV protease inhibitors ("PI"). In compliant drug-naïve patients, HAART is effective in reducing mortality and progression of HIV-1 to AIDS. However, these multidrug therapies do not eliminate HIV-1 and long-term treatment usually results in multidrug resistance. Development of new drug therapies to provide better HIV-1 treatment remains a priority.


SUMMARY OF THE INVENTION

The present invention provides the use of a pegylated interferon-alfa and a CCR5 antagonist for the preparation of a medicament for the treatment of HIV-1 infections in patients.






The present invention also provides the use of a pegylated interferon-alfa and a CCR5 antagonist for the preparation of a medicament for the treatment of HIV-1 infections in patients wherein the CCR5 antagonist is represented by the structural formula I or II or III or IV



3



X is -C(R¹³)₂-, -C(R¹³)(R¹⁹)-, -C(O)-, -O-, -NH-, -N((C₁-C₆)alkyl)-,




R is R⁶-phenyl, R⁶-pyridyl, R⁶-thiophenyl or R⁶-naphthyl;

R¹ is hydrogen, C₁-C₆ alkyl or C₂-C₆ alkenyl;

R² is R⁷, R⁸, R⁹-phenyl; R⁷, R⁸, R⁹-substituted 6-membered heteroaryl;

R⁷, R⁸, R⁹-substituted 6-membered heteroaryl N-oxide;

R¹⁰, R¹¹-substituted 5-membered heteroaryl; naphthyl; fluorenyl;



R³ is R⁶-phenyl, R⁶-heteroaryl or R⁶-naphthyl;

4


R⁴ is hydrogen, C₁-C₆ alkyl, fluoro-C₁-C₆ alkyl, cyclopropylmethyl, -CH₂CH₂OH, -CH₂CH₂-O-(C₁-C₆)alkyl, -CH₂C(O)-O-(C₁-C₆)alkyl,

-CH₂C(O)NH₂, -CH₂C(O)-NH(C₁-C₆)alkyl or -CH₂C(O)-N((C₁-C₆)alkyl)₂;

R⁵ and R¹¹ are independently selected from the group consisting of

hydrogen and (C₁-C₆)-alkyl;

R⁶ is 1 to 3 substituents independently selected from the group consisting of hydrogen, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, -CF₃, CF₃O-, CH₃C(O)-, -CN, CH₃SO₂-, CF₃SO₂-, R¹⁴-phenyl, R¹⁴-benzyl, CH₃C(=NOCH₃)₂-,



NHCO(C₁-C₆ alkyl), -NHSO₂(C₁-C₆ alkyl),

5-membered heteroaryl and —N—X , wherein X is -O-, -NH- or -N(CH₃)₂;

R⁷ and R⁸ are independently selected from the group consisting of (C₁-

C₆)alkyl, halogen, -NR²⁰R²¹, -OH, -CF₃, -OCH₃, -O-acyl, and -OCF₃;

R⁹ is R⁷, hydrogen, phenyl, -NO₂, -CN, -CH₂F, -CHF₂, -CHO,

-CH=NOR²⁰, pyridyl, pyridyl N-oxide, pyrimidinyl, pyrazinyl,

-N(R²⁰)CONR²¹R²², -NHCONH(chloro-(C₁-C₆)alkyl), -NHCONH((C₂-C₁₀)-

cycloalkyl)(C₁-C₆)alkyl), -NHCO(C₁-C₆)alkyl, -NHCOCF₃, -NHSO₂N((C₁-C₆)alkyl)₂, -

NHSO₂(C₁-C₆)alkyl, -N(SO₂CF₃)₂, -NHCO₂(C₁-C₆)alkyl, C₃-C₁₀ cycloalkyl, -SR²³, -

SOR²³, -SO₂R²³, -SO₂NH(C₁-C₆ alkyl), -OSO₂(C₁-C₆)alkyl, -OSO₂CF₃, hydroxy(C₁-

C₆)alkyl, -CON R²⁰R²¹, -CON(CH₂CH₂-O-CH₃)₂,

-OCONH(C₁-C₆)alkyl, -CO₂R²⁰, -Si(CH₃)₃ or -B(OC(CH₃)₃)₂;

R¹⁰ is (C₁-C₆)alkyl, -NH₂ or R¹²-phenyl;

R¹² is 1 to 3 substituents independently selected from the group consisting of hydrogen, (C₁-C₆) alkyl, -CF₃, -CO₂R²⁰, -CN, (C₁-C₆)alkoxy and halogen;

R¹³, R¹⁴, R¹⁵ and R¹⁶ are independently selected from the group consisting of hydrogen and (C₁-C₆)alkyl;

5

R¹⁷ and R¹⁸ are independently selected from the group consisting of hydrogen and C₁-C₆ alkyl, or R¹⁷ and R¹⁸ together are a C₂-C₅ alkylene group and with the carbon to which they are attached form a spiro ring of 3 to 6 carbon atoms;


- 5 R¹⁹ is R¹-phenyl, R²-heteroaryl, R³-naphthyl, C₂-C₆ cycloalkyl, (C₃-C₁₀)cycloalkyl/(C₁-C₆)alkyl or (C₁-C₆)alkoxy/(C₁-C₆)alkyl;

R²⁰, R²¹ and R²² are independently selected from the group consisting of H and C₁-C₆ alkyl; and

R²³ is C₁-C₆ alkyl or phenyl;



10

and wherein in the CCR5 antagonist compounds represented by the structural formula II:





or a pharmaceutically acceptable salt thereof, wherein

- 15 (1) X^a is -C(R¹³)₂-, -C(R¹³)(R¹⁹)-, -C(O)-, -O-, -NH-, -N((C₁-C₆)alkyl)-,



20



6




R^a is R^{8a}-phenyl, R^{8a}-pyridyl, R^{8a}-thiophenyl or R⁸-naphthyl;

R¹ is hydrogen, C₁-C₆ alkyl or C₂-C₆ alkenyl;

- 5 R² is R⁷, R⁸, R⁹-phenyl; R⁷, R⁸, R⁹-substituted 6-membered heteroaryl;

R⁷, R⁸, R⁹-substituted 6-membered heteroaryl N-oxide;

R¹⁰, R¹¹-substituted 5-membered heteroaryl; naphthyl; fluorenyl;



R³ is R¹⁰-phenyl, pyridyl, pyrimidyl, pyrazinyl or thiazolyl;

- 10 R⁴ is hydrogen, C₁-C₆ alkyl, fluoro-C₁-C₆ alkyl, cyclopropyl/methyl,

-CH₂CH₂OH, -CH₂CH₂-O-(C₁-C₆)alkyl, -CH₂C(O)-O-(C₁-C₆)alkyl,


-CH₂C(O)NH₂, -CH₂C(O)-NH(C₁-C₆)alkyl or -CH₂C(O)-N((C₁-C₆)alkyl)₂;

R⁵ and R¹¹ are independently selected from the group consisting of

hydrogen and (C₁-C₆)-alkyl;

15

R^{6a} is 1 to 3 substituents independently selected from the group consisting of hydrogen, halogen, -CF₃, CF₃O-, -CN, -CF₃SO₂, R¹²-phenyl,

-NHCOOCF₃, 5-membered heteroaryl and , wherein X is -O-, -NH- or -N(CH₃)₂;

R⁶ is independently selected from the group consisting of R^{6a} and

20 CH₃SO₂-;

20

R⁷ and R⁸ are independently selected from the group consisting of (C₁-C₆)alkyl, halogen, -NR²⁰R²¹, -OH, -CF₃, -OCH₃, -O-acyl, and -OCF₃;

R⁹ is R¹, hydrogen, phenyl, -NO₂, -CN, -CH₂F, -CHF₂, -CHO,

-CH=NO²⁰, pyridyl, pyridyl N-oxide, pyrimidinyl, pyrazinyl,

- 25 -N(R²⁰)CONR²¹R²², -NHCONH(chloro-(C₁-C₆)alkyl), -NHCONH((C₃-C₁₀)-

7

cycloalkyl(C₁-C₆alkyl), -NHCO(C₁-C₆alkyl), -NHCOCF₃, -NHSO₂N((C₁-C₆alkyl)₂), -NHSO₂(C₁-C₆alkyl), -N(SO₂CF₃)₂, -NHCO₂(C₁-C₆alkyl), C₃C₁₀ cycloalkyl, -SR²³, -SOR²³, -SO₂R²³, -SO₂NH(C₁-C₆alkyl), -OSO₂(C₁-C₆alkyl), -OSO₂CF₃, hydroxy(C₁-C₆alkyl), -CON R²⁰R²¹, -CON(CH₂CH₂-O-CH₃)₂,

5 -OCONH(C₁-C₆alkyl), -CO₂R²⁰, -Si(CH₃)₃ or -B(OC(CH₃)₂)₂;

R¹⁰ is (C₁-C₆alkyl), -NH₂ or R¹²-phenyl;

R¹² is 1 to 3 substituents independently selected from the group consisting of hydrogen, (C₁-C₆alkyl), -CF₃, -CO₂R₂₀, -CN, (C₁-C₆alkoxy and halogen;

R¹³, R¹⁴, R¹⁵ and R¹⁶ are independently selected from the group

10 consisting of hydrogen and (C₁-C₆alkyl);

R¹⁷ and R¹⁸ are independently selected from the group consisting of

hydrogen and C₁-C₆alkyl, or R¹⁷ and R¹⁸ together are a C₂-C₅ alkylene group and with the carbon to which they are attached form a spiro ring of 3 to 6 carbon

atoms;

15 R¹⁹ is R²-phenyl, R²-heteroaryl, R²-naphthyl, C₂-C₁₀ cycloalkyl, (C₂-


C₁₀cycloalkyl)(C₁-C₆alkyl) or (C₁-C₆alkoxy)(C₁-C₆alkyl);

R²⁰, R²¹ and R²² are independently selected from the group consisting of H and C₁-C₆alkyl; and


R²³ is C₁-C₆alkyl or phenyl; or

20 (2):



X^a is -C(R¹³)(R¹⁹), -C(O)-, -O-, -NH-, -N((C₁-C₆alkyl))-,



25



8



5

R^a is R^{8b}-phenyl, R^{8b}-pyridyl or R^{8b}-thiophenyl;


R^{4a} is fluoro-C₁-C₆alkyl, cyclopropylmethyl, -CH₂CH₂OH,

-CH₂CH₂-O-(C₁-C₆alkyl), -CH₂C(O)-O-(C₁-C₆alkyl), -CH₂C(O)NH₂, -CH₂C(O)-NH-(C₁-C₆alkyl) or -CH₂C(O)-N((C₁-C₆alkyl)₂;

R^{8b} is CH₃SO₂-; and

10 R¹, R², R³, R⁵, R¹⁴, R¹⁵, R¹⁶ and R¹⁹ are as defined in II(1);

and wherein in the CCR5 antagonist compounds represented by the structural formula III:



III


15

R is R⁸-phenyl, R⁸-pyridyl, R⁸-thiophenyl or R⁸-naphthyl;

R¹ is hydrogen or C₁-C₆alkyl;

R² is R⁹, R¹⁰, R¹¹-phenyl; R⁹, R¹⁰, R¹¹-substituted 6-membered heteroaryl; R⁹, R¹⁰, R¹¹-substituted 5-membered heteroaryl; naphthyl; fluorenyl;

20



9


R³ is hydrogen, C₁-C₆ alkyl, (C₁-C₆)alkoxy(C₁-C₆)alkyl, C₃-C₁₀ cycloalkyl, C₃-C₁₀ cycloalkyl(C₁-C₆)alkyl, R⁸-phenyl, R⁸-phenyl(C₁-C₆)alkyl, R⁸-naphthyl, R⁸-naphthyl(C₁-C₆)alkyl, R⁸-heteroaryl or R⁸-heteroaryl(C₁-C₆)alkyl;

5 R⁴, R⁵, R⁷ and R¹³ are independently selected from the group consisting of hydrogen and (C₁-C₆)-alkyl;

R⁶ is hydrogen, C₁-C₆ alkyl or C₂-C₆ alkenyl;

R⁸ is 1 to 3 substituents independently selected from the group consisting of hydrogen, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, -CF₃, CF₃O-, CH₃C(O)-, -CN, CH₃SO₂-, CF₃SO₂-, R¹⁴-phenyl, R¹⁴-benzyl, CH₃C(=NOCH₃),


10 CH₃C(=NOCH₂CH₃),



, -NH₂, -NHCOCF₃,

-NHCONH(C₁-C₆ alkyl), -NHCO(C₁-C₆ alkyl), -NHSO₂(C₁-C₆ alkyl),



5-membered heterocaryl and , wherein X is -O-, -NH- or -N(CH₃);

R⁹ and R¹⁰ are independently selected from the group consisting of (C₁-C₆)alkyl, halogen, -NR¹⁷R¹⁸, -OH, -CF₃, -OCH₃, -O-acyl, -OCF₃ and

15 -Si(CH₃)₃;

R¹¹ is R⁸, hydrogen, phenyl, -NO₂, -CN, -CH₂F, -CHF₃, -CHO, -CH=NOR¹⁷, pyridyl, pyridyl N-oxide, pyrimidinyl, pyrazinyl,

-N(R¹⁷)CONR¹⁸R¹⁹, -NHCONH(chloro-(C₁-C₆)alkyl), -NHCONH((C₃-

20 C₁)cycloalkyl(C₁-C₆)alkyl), -NHCO(C₁-C₆)alkyl, -NHCOCF₃, -NHSO₂N((C₁-C₆)alkyl)₂, -NHSO₂(C₁-C₆)alkyl, -N(SO₂CF₃)₂, -NHCO₂(C₁-C₆)alkyl, C₃-C₁₀ cycloalkyl, -SR²⁰, -SOR²⁰, -SO₂R²⁰, -SO₂NH(C₁-C₆ alkyl), -OSO₂(C₁-C₆)alkyl, -OSO₂CF₃, hydroxy(C₁-C₆)alkyl, -CON R¹⁷R¹⁸, -CON(CH₂CH₂O-CH₃)₂, -OCONH(C₁-C₆)alkyl, -CO₂R¹⁷, -Si(CH₃)₃ or -B(OC(CH₃)₂)₂;

R¹² is (C₁-C₆)alkyl, -NH₂ or R¹⁴-phenyl;

25 R¹⁴ is 1 to 3 substituents independently selected from the group consisting of hydrogen, (C₁-C₆) alkyl, -CF₃, -CO₂R¹⁷, -CN, (C₁-C₆)alkoxy and halogen;


10

R¹⁵ and R¹⁶ are independently selected from the group consisting of hydrogen and C₁-C₆ alkyl, or R¹⁵ and R¹⁶ together are a C₂-C₅ alkylene group and with the carbon to which they are attached form a spiro ring of 3 to 6 carbon atoms;

5 R¹⁷, R¹⁸ and R¹⁹ are independently selected from the group consisting of H and C₁-C₆ alkyl; and

R²⁰ is C₁-C₆ alkyl or phenyl;

10 and wherein in the CCR5 antagonist compounds represented by the structural formula IV:




IV

or a pharmaceutically acceptable salt thereof, wherein

(1) R^a is R^{8a}-phenyl, R^{8b}-pyridyl, R^{8b}-thiophenyl or R⁸-naphthyl;

R¹ is hydrogen or C₁-C₆ alkyl;

15 R² is R⁹, R¹⁰, R¹¹-phenyl; R⁹, R¹⁰, R¹¹-substituted 6-membered heteroaryl; R⁹, R¹⁰, R¹¹-substituted 6-membered heteroaryl N-oxide; R¹², R¹³-substituted 5-membered heteroaryl; naphthyl; fluorenyl;



;


20 R³ is hydrogen, C₁-C₆ alkyl, (C₁-C₆)alkoxy(C₁-C₆)alkyl, C₃-C₁₀ cycloalkyl, C₃-C₁₀ cycloalkyl(C₁-C₆)alkyl, R⁸-phenyl, R⁸-phenyl(C₁-C₆)alkyl, R⁸-naphthyl, R⁸-naphthyl(C₁-C₆)alkyl, R⁸-heteroaryl or R⁸-heteroaryl(C₁-C₆)alkyl;

R⁴, R⁵, R⁷ and R¹³ are independently selected from the group consisting of hydrogen and (C₁-C₆)-alkyl;


R⁶ is hydrogen, C₁-C₆ alkyl or C₂-C₆ alkenyl;

11


R⁸ is 1 to 3 substituents independently selected from the group consisting of hydrogen, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, -CF₃, CF₃O-, CH₃C(O)-, -CN, CH₃SO₂-, CF₃SO₂-, R¹⁴-phenyl, R¹⁴-benzyl, CH₃C(=NOCH₃),



5 -NHCONH(C1-C6 alkyl), -NHCO(C1-C6 alkyl), -NH₂SO₂(C1-C6 alkyl),



R^{8e} is 1 to 3 substituents independently selected from the group consisting of hydrogen, halogen, -CF₃, CF₃O-, -CN, CF₃SO₂-, R¹⁴-phenyl, -NHCOCF₃, 5-



membered heterocaryl and , wherein X is as defined above;

10 R8b is 1 to 3 substituents independently selected from the group consisting of hydrogen, halogen, -CF₃, CF₃O-, CH₃C(O)-, -CN, CF₃SO₂-, R¹⁴-benzyl,



-NHCOCF₃, 5-membered heterocaryl and , wherein X is as defined above;

R⁹ and R¹⁰ are independently selected from the group consisting of (C₁-

15 C_6 /alkyl, halogen, $-NR''R'''$, $-OH$, $-CF_3$, $-OCH_3$, $-O$ -acyl, $-OCF_3$ and

$$-\text{Si}(\text{CH}_3)_3;$$

R¹¹ is R⁹, hydrogen, phenyl, -NO₂, -CN, -CH₂F, -CHF₂, -CHO,

-CH=NOR¹⁷, pyridyl, pyridyl N-oxide, pyrimidinyl, pyrazinyl,

$-\text{N}(\text{R}^{17})\text{CONR}^{18}\text{R}^{18}$, $-\text{NHCONH}(\text{chloro}-(\text{C}_1-\text{C}_6\text{alkyl}), -\text{NHCONH}((\text{C}_3-$

20 C₉cycloalkyl(C₁-C₆alkyl), -NHCO(C₁-C₆alkyl), -NHCOCF₃, -NHSO₂N((C₁-C₆alkyl)₂), -NHSO₂(C₁-C₆alkyl), -N(SO₂CF₃)₂, -NHCO₂(C₁-C₆alkyl), C₃C₁₀cycloalkyl, -SR²⁰, -SOR²⁰, -SO₂R²⁰, -SO₂NH(C₁-C₆alkyl), -OSO₂(C₁-C₆alkyl),

12

$$\text{OSO}_2\text{CF}_3, \text{hydroxy}(\text{C}_1\text{-C}_8\text{)alkyl}, -\text{CON R}^{17}\text{R}^{18}, -\text{CON}(\text{CH}_2\text{CH}_2\text{CH}_2\text{-O-CH}_3)_2$$
$$-\text{OCONH}(\text{C}_1\text{-C}_6)\text{alkyl}, -\text{CO}_2\text{R}^{17}, -\text{Si}(\text{CH}_3)_3 \text{ or } -\text{B}(\text{OC}(\text{CH}_3)_2)_2$$

R¹² is (C₁-C₆)alkyl, -NH₂ or R¹⁴-phenyl;

R¹⁴ is 1 to 3 substituents independently selected from the group consisting of hydrogen, (C₁-C₆) alkyl, -CF₃, -CO₂R¹⁷, -CN, (C₁-C₆)alkoxy and halogen;


R¹⁵ and R¹⁶ are independently selected from the group consisting of

hydrogen and C₁-C₆ alkyl, or R¹⁵ and R¹⁶ together are a C₂-C₅ alkylene group and with the carbon to which they are attached form a spiro ring of 3 to 6 carbon atoms;

¹⁰ R¹⁷, R¹⁸ and R¹⁹ are independently selected from the group consisting of H, C₁-C₆ alkyl, and C₁-C₆ alkyl; and

R^{20} is C₁-C₈ alkyl or phenyl; or

(2) R^a is R⁸-phenyl, R⁸-pyridyl or R⁸-thiophenyl;



R² is fluorenyl, diphenylmethyl,

15 and $R^1, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17},$
 R^{18}, R^{19} and R^{20} are as defined in IV(1).

Preferred are compounds of formula I wherein R is R⁶-phenyl, especially


wherein R⁶ is a single substituent, and especially wherein the R⁶ substituent is in the 4-position. Also preferred are compounds of formula I wherein R¹³, R¹⁴, R¹⁵

and R¹⁵ are each hydrogen or methyl, especially hydrogen. Also preferred are compounds of formula I wherein X is -CHOR³, -C(R¹³)(R¹⁴)- or -C(=NOR⁴); a preferred definition for R³ is pyridyl, especially 2-pyridyl, a preferred definition for

R⁴ is (C₁-C₆)alkyl, especially methyl, ethyl or isopropyl, a preferred definition for **R⁴** is hydrogen, and a preferred definition for **R⁴** is **R⁴**-phenyl. For compounds of formula I, **R¹** is preferably (C₁-C₆)alkyl, especially methyl.

13

In compounds of formula I, R² is preferably R⁷, R⁸, R⁹-phenyl, R⁷, R⁸, R⁹-pyridyl or an N-oxide thereof, or R⁷, R⁸, R⁹-pyrimidyl. When R² is pyridyl, it is preferably 3- or 4-pyridyl, and when pyrimidyl, it is preferably 5-pyrimidyl. The R⁷ and R⁸ substituents are preferably attached to carbon ring members adjacent to the carbon joining the ring to the rest of the molecule and the R⁹ substituent can be attached to any of the remaining unsubstituted carbon ring members, for example as shown in the following structures:



Preferred R⁷ and R⁸ substituents are: (C₁-C₆alkyl), especially methyl; halogen, especially chloro; and -NH₂. A preferred R⁹ substituent is hydrogen.

Preferred are compounds of formula II(1) wherein R^a is R^{8b}-phenyl, especially wherein R^{8b} is a single substituent, and especially wherein the R^{8b} substituent is in the 4-position. Also preferred are compounds of formula II(1) wherein X^a is -CHOR³, -C(R³)(R^{3'})- or -C(=NOR⁴); a preferred definition for R³ is pyridyl, especially 2-pyridyl, a preferred definition for R⁴ is (C₁-C₆alkyl), especially methyl, ethyl or isopropyl, a preferred definition for R^{3'} is hydrogen, and a preferred definition for R^{3'} is R⁴-phenyl. For compounds of formula II(1), R¹ is preferably (C₁-C₆alkyl), especially methyl. Also for compounds of formula II(1), R¹⁴, R¹⁵ and R¹⁶ are preferably hydrogen.

Preferred are compounds of formula II(2) wherein R^a is R^{8b}-phenyl, especially wherein R^{8b} is a single substituent, and especially wherein the R^{8b} substituent is in the 4-position. Also preferred are compounds of formula II(2) wherein X^a is -CHOR³, -C(R³)(R^{3'})- or -C(=NOR⁴); a preferred definition for R³ is pyridyl, especially 2-pyridyl, preferred definitions for R⁴ are cyclopropylmethyl and trifluoroethyl, a preferred definition for R^{3'} is hydrogen, and a preferred definition for R^{3'} is R⁴-phenyl. For compounds of formula II(2), R¹ is preferably


14

(C₁-C₆alkyl), especially methyl. Also for compounds of formula II(2), R¹⁴, R¹⁵ and R¹⁶ are preferably hydrogen.

In compounds of formula II(1) and (2), R² is preferably R⁷, R⁸, R⁹-phenyl; R⁷, R⁸, R⁹-pyridyl or an N-oxide thereof, or R⁷, R⁸, R⁹-pyrimidyl. When R² is pyridyl, it is preferably 3- or 4-pyridyl, and when pyrimidyl, it is preferably 5-pyrimidyl. The R⁷ and R⁸ substituents are preferably attached to carbon ring members adjacent to the carbon joining the ring to the rest of the molecule and the R⁹ substituent can be attached to any of the remaining unsubstituted carbon ring members as shown above for compounds of formula I. Preferred R⁷ and R⁸ substituents for compounds of formula II are: (C₁-C₆alkyl), especially methyl; halogen, especially chloro; and -NH₂; a preferred R⁹ substituent is hydrogen.

Preferred are compounds of formula III wherein R¹ is R⁸-phenyl or R⁸-naphthyl, especially wherein R⁸ is a single substituent, and especially wherein the R⁸ substituent is in the 4-position. For R⁸-phenyl, preferred R⁸ substituents are -CF₃, -OCF₃, CH₃SO₂-, CH₃CO-, CH₃C(=NOCH₃)-, Br and I. For R⁸-naphthyl, R⁸ is preferably C₁-C₆ alkoxy. Also preferred are compounds of formula III wherein R³ is hydrogen, (C₁-C₆) alkyl, R⁸-phenyl, R⁸-benzyl or R⁸-pyridyl; more preferred definitions for R³ are methyl, ethyl, phenyl, benzyl and pyridyl. R¹ is preferably hydrogen. For compounds of formula III, R⁶ is preferably hydrogen or methyl, especially methyl. R⁴ is preferably methyl; R⁵ and R⁷ are each preferably hydrogen.

In compounds of formula III, R² is preferably R⁹, R¹⁰, R¹¹-phenyl, R⁹, R¹⁰, R¹¹-pyridyl or an N-oxide thereof, or R⁹, R¹⁰, R¹¹-pyrimidyl. When R² is pyridyl, it is preferably 3- or 4-pyridyl, and when pyrimidyl, it is preferably 5-pyrimidyl. The R⁹ and R¹⁰ substituents are preferably attached to carbon ring members adjacent to the carbon joining the ring to the rest of the molecule and the R¹¹ substituent can be attached to any of the remaining unsubstituted carbon ring members, for example as shown in the following structures:



15


Preferred R⁹ and R¹⁰ substituents are: (C₁-C₆)alkyl, especially methyl;

halogen, especially chloro or bromo, -OH and -NH₂. When R² is phenyl, R¹¹ is

preferably hydrogen or -OH; when R² is pyridyl, R¹¹ is preferably hydrogen; and

when R² is pyrimidyl, R¹¹ is preferably hydrogen, methyl or phenyl. Examples of

particularly preferred R² groups are as follows:



10

Preferred compounds of formula IV are those defined in (1).

More preferred are those of formula IV(1) wherein R^{8a} is R^{8a}-phenyl or R^{8a}-

naphthyl, wherein R^{8a} is -CF₃, CF₃O- or halogen and R⁸ is C₁-C₆ alkoxyl. The

R^{8a} or R⁸ substituent is preferably a single substituent; it is especially preferred

that the R^{8a} or R⁸ substituent is in the 4-position. Also preferred are compounds

of formula IV(1) wherein R³ is hydrogen, (C₁-C₆) alkyl, R⁸-phenyl, R⁸-benzyl or

R⁸-pyridyl; more preferred definitions for R³ are methyl, ethyl, phenyl, benzyl and

pyridyl. R¹ is preferably hydrogen. For compounds of formula IV(1), R⁶ is

preferably hydrogen or methyl, especially methyl. R⁴ is preferably methyl, R⁵ and R⁷ are each preferably hydrogen.

R² in formula IV(1) is preferably as defined for formula III, i.e., R⁹, R⁹, R¹⁰,

R¹¹-phenyl, R⁹, R¹⁰, R¹¹-pyridyl or an N-oxide thereof, or R⁹, R¹⁰, R¹¹-pyrimidyl,

wherein the R⁹, R¹⁰, R¹¹-substitution is as defined above for preferred

compounds of formula III.

The present invention also provides the use of a pegylated interferon- α , a

CCRS antagonist, ribavirin and HAART for the preparation of a medicament for

the treatment of patients co-infected with HIV-1 and HCV. The treatment


comprises administering a therapeutically effective amount of pegylated

interferon- α in association with a therapeutically effective amount of ribavirin




and a therapeutically effective amount of HAART and a therapeutically effective

amount of a CCR5 antagonist represented by the structural formula I or II or III or

IV:



15



or a pharmaceutical salt of a compound of the formula I or II or III or IV;

sufficient to lower HIV-1-RNA and HCV-RNA plasma levels.

wherein X, R, and R¹ to R¹⁶ are as defined in formula I; and

wherein X, R⁺ and R¹ to R¹⁶ are as defined in formula II(1) and II(2) and


16

17

wherein R₁ and R₁' to R₁₇' are as defined in formula II; and wherein R^a, and R¹ to R¹⁷ are as defined in formula IV.

The present invention also provides the use of pegylated interferon- α , HAART and a CCR5 antagonist for the preparation of a medicament for the treatment of pediatric patients infected with HIV-1. The treatment comprises administering a therapeutically effective amount of pegylated interferon- α in association with a therapeutically effective amount of HAART and a therapeutically effective amount of a CCR5 antagonist represented by the structural formula I or II or III or IV.

10



or a pharmaceutical salt of compounds I or II or III or IV; sufficient to lower HIV-1-RNA plasma levels.

wherein X, R₁ and R₁' to R₁₆ are as defined in I; and wherein X^a, R^a and R¹ to R₁₆ are as defined in II(1) and II(2); wherein R₁ and R₁' to R₁₇' are as defined in formula II; and wherein R^a, and R¹ to R₁₇' are as defined in formula IV.

20

18

DETAILED DESCRIPTION

As used herein, the following terms are used as defined below unless otherwise indicated.

5 Alkyl (including the alkyl portions of alkoxy, alkylamino and dialkylamino) represents straight and branched carbon chains and contains from one to six carbon atoms.

Alkenyl represents C₂-C₆ carbon chains having one or two unsaturated bonds, provided that two unsaturated bonds are not adjacent to each other.

10 Substituted phenyl means that the phenyl group can be substituted at any available position on the phenyl ring.

Acyl means a radical of a carboxylic acid having the formula alkyl-C(O)-, aryl-C(O)-, aralkyl-C(O)-, (C₃-C₇)cycloalkyl-C(O)-, (C₃-C₇)cycloalkyl-(C₁-C₆)alkyl-C(O)-, and heteroaryl-C(O)-, wherein alkyl and heteroaryl are as defined herein; aryl is R¹²-phenyl or R¹²-naphthyl; and aralkyl is aryl-(C₁-C₆)alkyl, wherein aryl is as defined above.

15

Heteroaryl represents cyclic aromatic groups of 5 or 6 atoms or bicyclic groups of 11 to 12 atoms having one or two heteroatoms independently selected from O, S or N, said heteroatom(s) interrupting a carbocyclic ring structure and having a sufficient number of delocalized pi electrons to provide aromatic character. For 6-membered heteroaryl rings, carbon atoms can be substituted by R⁷, R⁸ or R⁹ groups. Nitrogen atoms can form an N-oxide. All regioisomers are contemplated, e.g., 2-pyridyl, 3-pyridyl and 4-pyridyl. Typical 6-membered heteroaryl groups are pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl and the N-oxides thereof. For 5-membered heteroaryl rings, carbon atoms can be substituted by R¹⁰ or R¹¹ groups. Typical 5-membered heteroaryl rings are furyl, thienyl, pyrrolyl, thiazolyl, isothiazolyl, imidazolyl, pyrazolyl and isoxazolyl. 5-Membered rings having one heteroatom can be joined through the 2- or 3- position; 5-membered rings having two heteroatoms are preferably joined through the 4-position. Bicyclic groups typically are benzo-fused ring systems derived from the heteroaryl groups named above, e.g. quinolyl, phthalazinyl, quinoxalinyl, benzofuranyl, benzothienyl and indolyl.

30

Preferred points of substitution for 6-membered heteroaryl rings at R² are described above. When R² is a 5-membered heteroaryl group, the R¹⁰ and R¹¹ substituents are preferably attached to carbon ring members adjacent to the carbon joining the ring to the rest of the molecule, and R¹¹ is preferably alkyl;

35

19

however, if a heteroatom is adjacent to the carbon joining the ring to the rest of the molecule (i.e., as in 2-pyrrolyl), R¹⁰ is preferably attached to a carbon ring member adjacent to the carbon joining the ring to the rest of the molecule.

5 Halogen represents fluoro, chloro, bromo and iodo.

Fluoro(C₁-C₆)alkyl represents a straight or branched alkyl chain substituted by 1 to 5 fluoro atoms, which can be attached to the same or different carbon atoms, e.g., -CH₂F, -CHF₂, -CF₃, F₃CCH₂- and -CF₂CF₃.

10 The present method of treating patients having HIV-1 infections comprises administering a therapeutically effective amount of pegylated interferon- α and a therapeutically effective amount of a CCR5 antagonist compound represented by structural formula I or II or III or IV as a combination therapy or in association with a therapeutically effective amount of at least one of ribavirin, interleukin-2("IL-2"), interleukin-12("IL-12"), and pentaluside alone or in combination with an anti-HIV-1 therapy, especially, HAART in accordance with good clinical practice to minimize HIV-1-RNA plasma levels. See for example A-M. Vandamme et al., in *Antiviral Chemistry & Chemotherapy*, 9:187-203 (1998) and "Drugs for HIV Infection" in *The Medical Letter* Vol. 39 (Issue 1015) December 5, 1997, pages 111-116. In a preferred aspect of the present invention, the combination of a pegylated interferon α and a CCR5 antagonist of formulas I to IV is administered to a patient infected with HIV-1, or co-infected with HIV-1 and HCV, in association with ribavirin and HAART. It is a special feature of the present invention that each of pegylated interferon α , the CCR5 antagonists of formulas I to IV and the components of HAART has a different mechanism of action in treating HIV-1.

30 It is another special feature of the present invention that the pegylated interferon α and the CCR5 antagonists of formulas I to IV do not cause cross-resistance with each other or with the components of HAART. The initiation of the administration of a therapeutically effective amount of the combination of a pegylated interferon α , ribavirin, and a CCR5 antagonist compound represented by structural formula I or II or III or IV and HAART may occur before,

20

after or concurrently with administering a therapeutically effective amount of combination of a pegylated interferon- α and a CCR5 antagonist compound represented by structural formula I or II or III or IV in accordance with the present invention. In an embodiment of the present invention, the method of treating patients having HIV-1 infections comprises two treatment time periods. In the first treatment time period, a combination of a therapeutically effective amount of pegylated interferon- α and a CCR5 antagonist compound represented by structural formula I or II or III or IV is administered for a first treatment time period sufficient to lower HIV-1-RNA plasma levels, preferably by a power of 10, more preferably by at least two powers of ten, i.e., at least 10², lower than the initial HIV-1-RNA plasma level. In the second treatment time period, the method entails continuing the administration of a therapeutically effective amount of a combination of pegylated interferon- α in association a CCR5 antagonist compound represented by structural formula I or II or III or IV and a therapeutically effective amount of HAART in accordance with good clinical practice to minimize HIV-1-RNA plasma levels. A-M. Vandamme et al., *Antiviral Chemistry & Chemotherapy*, 9:187-203 (1998) disclose current clinical treatments of HIV-1 infections, including when to start multiring therapy and which drugs to combine. The triple drug therapy may include two NRTIs and one PI, but there are many issues to be considered in the choice of the precise HAART for any patient. See for example, Tables 1 & 2 and Figure 2 in A-M. Vandamme et al., listed hereinabove.

25 The terms "CCR5 antagonist compound" and "CCR5 antagonists" as used herein mean any compound that interferes with the interaction between the viral receptor CCR5 and HIV-1 to block entry of HIV-1 into the cell. Assays, e.g., the CCR5 Membrane Binding Assay, the HIV-1 Entry and the HIV-1 Entry Replication Assays, inter alia, are presented herein after to identify a compound as a CCR5 antagonist and to determine its CCR5 antagonist activity.

30

- 5 The term "patients having HIV-1 infections" as used herein means any patient -including a pediatric patient-having HIV-1 infection and includes treatment-naïve patients and treatment-experienced patients having the HIV-1 infection as well as treatment-naïve patients and treatment-experienced patients co-infected with the HIV-1 and hepatitis C virus ("HCV").

10 The term "pediatric patient" as used herein means a patient below the age of 17, and normally includes those from birth to 16 years of age.

- 15 The term "treatment-naïve patients" as used herein means patients having HIV-1 or co-infected with the HIV-1 and HCV who have never been treated with any anti-retroviral drugs, e.g., NRTI, NNRTI, PI or any interferon, including but not limited to interferon- α , or pegylated interferon α .

- 20 The term "treatment-experienced" patients as used herein means those patients having HIV-1 or co-infected with the HIV-1 and HCV who have initiated some form of anti HIV therapy including, but not limited to HAART or some form of anti-HCV therapy, including but not limited to interferon- α , pegylated interferon α or ribavirin.

- 25 The term "patients having hepatitis C infections" as used herein means any patient-including a pediatric patient- having hepatitis C and includes treatment-naïve patients having hepatitis C infections and treatment-experienced patients having hepatitis C infections as well as those pediatric, treatment-naïve and treatment-experienced patients having chronic hepatitis C infections.

- 30 These patients having hepatitis C include those who are infected with multiple HCV genotypes including type 1 as well as those infected with, e.g., HCV genotypes 2, 3, 4, 5 and/or 6 and other possible HCV genotypes.

- 5 The term "treatment-naïve patients having hepatitis C infections" as used herein means patients with hepatitis C who have never been treated with ribavirin or any interferon, including but not limited to interferon- α , or pegylated interferon α .

- 10 The term "treatment-experienced patients having hepatitis C infections" as used herein means patients with hepatitis C who have been treated with ribavirin or any interferon, including but not limited to interferon- α , or pegylated interferon α , including relapsers and non-responder.

- 15 The term "relapsers" as used herein means treatment-experienced patients with hepatitis C who have relapsed after initial response to previous treatment with interferon alone, or in combination with ribavirin.

- 20 The term "non-responders" as used herein means treatment-experienced patients with hepatitis C who have not responded to prior treatment with any interferon alone, or in combination with ribavirin.

- 25 When the pegylated interferon- α administered is a pegylated interferon α -2b, the therapeutically effective amount of pegylated interferon α -2b administered during the treatment in accordance with the present invention, including in first and second treatment time periods, is in the range of about 0.1 to 9.0 micrograms per kilogram of pegylated interferon α -2b administered per week, in single or divided doses, preferably once a week (QW) or twice a week (BIW), preferably in the range of about 0.1 to about 9.0 micrograms per kilogram of pegylated interferon α -2b administered once a week (QW) or in the range of about 0.05 to about 4.5 micrograms per kilogram of pegylated interferon α -2b administered twice a week (BIW), or is in the range of about 0.5 to about 3.0 micrograms per kilogram of pegylated interferon α -2b administered per week, preferably in the range of about 0.5 to about 3.0 micrograms per kilogram of pegylated interferon α -2b administered once a week (QW) or in the range of

23

about 0.25 to about 1.5 micrograms per kilogram of pegylated interferon alfa-2b administered twice a week, or is in the range of about 0.75 to about 1.5 micrograms per kilogram of pegylated interferon alfa-2b administered per week, most preferably is in the range of about 0.75 to about 1.5 micrograms per kilogram of pegylated interferon alfa-2b administered once a week or about 0.375 to about 0.75 micrograms per kilogram of pegylated interferon alfa-2b administered twice a week.

10 When the pegylated interferon-alfa administered to pediatric patients is a pegylated interferon alfa-2b, the therapeutically effective amount of pegylated interferon alfa-2b administered during the treatment in accordance with the present invention, including in first and second treatment time periods is in the range of about 0.1 to 9.0 micrograms per kilogram of pegylated interferon alfa-2b administered per week, in single or divided doses, preferably once a week (QW) or twice a week (BIW), more preferably about 0.1 to about 9.0 micrograms per kilogram of pegylated interferon alfa-2b administered once a week (QW), or about 0.05 to about 4.5 micrograms per kilogram of pegylated interferon alfa-2b administered per week, in single or divided doses, preferably once a week (QW) or twice a week (BIW), more preferably about 0.05 to about 4.5 micrograms per kilogram of pegylated interferon alfa-2b administered once a week, or preferably about 0.75 to about 3.0 micrograms per kilogram of pegylated interferon alfa-2b administered in single or divided doses, preferably once a week (QW) or twice a week (BIW), more preferably about 0.75 to about 3.0 micrograms per kilogram of pegylated interferon alfa-2b administered once a week or about 0.375 to about 1.5 micrograms per kilogram of pegylated interferon alfa-2b administered twice a week, and most preferably about 2.25 to about 2.6 micrograms per kilogram of pegylated interferon alfa-2b administered once a week or about 1.1 to about 1.3 micrograms per kilogram of pegylated interferon alfa-2b administered twice a week (BIW).

24

When the pegylated interferon-alfa administered is a pegylated interferon alfa-2a, the therapeutically effective amount of pegylated interferon alfa-2a administered during the treatment in accordance with the present invention, including in first and second treatment time periods, is in the range of about 50 micrograms to about 500 micrograms per week, preferably about 150 micrograms to about 250 micrograms per week, or preferably about 180 micrograms to about 250 micrograms per week or preferably about 150 micrograms to about 180 micrograms per week or most preferably about 180 micrograms per week or alternatively the effective amount is in the range of about 50 micrograms to about 500 micrograms once a week ("QW"), preferably about 150 micrograms to about 250 micrograms QW, or preferably about 180 micrograms to about 250 micrograms QW or preferably about 150 micrograms to about 180 micrograms QW or most preferably about 180 micrograms QW or alternatively the effective amount is in the range of about 25 micrograms to about 250 micrograms twice a week ("BIW"), preferably about 75 micrograms to about 125 micrograms BIW, preferably about 75 micrograms to about 125 micrograms BIW, or preferably about 75 micrograms to about 90 micrograms BIW, or most preferably about 90 micrograms BIW.

20 When the pegylated interferon-alfa administered to a pediatric patient is a pegylated interferon alfa-2a, the therapeutically effective amount of pegylated interferon alfa-2a administered during the treatment in accordance with the present invention, including in first treatment time period is in the range of about 50 micrograms to about 500 micrograms once a week ("QW"), preferably about 300 micrograms to about 375 micrograms QW or the therapeutically effective amount of pegylated interferon alfa-2a administered to a pediatric patient is in the range of about 50 micrograms to about 250 micrograms twice a week, preferably about 150 micrograms to about 190 micrograms once a week.

30 Ribavirin is administered to the patient in association with pegylated interferon-alfa, that is, before, after or concurrently with the administration of the

25

pegylated interferon alfa. The pegylated interferon-alfa dose is preferably administered during the same period of time that the patient receives doses of ribavirin. The amount of ribavirin administered concurrently with the pegylated interferon-alfa is from about 400 to about 1600 mg per day, preferably about 600 to about 1200 mg/day or about 800 to about 1200 mg day and most preferably about 1000 to about 1200 mg/kg a day. The pegylated interferon-alfa dose is also preferably administered to the pediatric patient during the same period of time that such patient receives doses of ribavirin. The amount of ribavirin administered to the pediatric patient concurrently with the pegylated interferon-alfa is from about 8 to about 15 mg per kilogram per day, preferably about 8, 12 or 15 mg per kilogram per day, in divided doses.

Pegylated interferon-alfa formulations are not effective when administered orally, so the preferred method of administering the pegylated interferon-alfa is parenterally. Preferably by subcutaneous, IV, or IM, injection. Ribavirin may be administered orally in capsule, tablet or liquid form in association with the parenteral administration of pegylated interferon-alfa. Of course, other types of administration of both medications, as they become available are contemplated, such as by nasal spray, transdermally, by suppository, by sustained release dosage form, and by pulmonary inhalation. Any form of administration will work so long as the proper dosages are delivered without destroying the active ingredient.

The term "nucleoside and nucleotide reverse transcriptase inhibitors" ("NRTI"s) as used herein means nucleosides and nucleotides and analogues thereof that inhibit the activity of HIV-1 reverse transcriptase, the enzyme which catalyzes the conversion of viral genomic HIV-1 RNA into proviral HIV-1 DNA.

Typical suitable NRTIs include zidovudine (AZT) available under the RETROVIR tradename from Glaxo-Wellcome Inc., Research Triangle, NC 27709; didanosine (ddI) available under the VIDEX tradename from Bristol-Myers

26

Squibb Co., Princeton, NJ 08543; zalcitabine (ddC) available under the HIVID tradename from Roche Pharmaceuticals, Nutley, NJ 07110; stavudine (d4T) available under the ZERIT trademark from Bristol-Myers Squibb Co., Princeton, NJ 08543; lamivudine (3TC) available under the EPIVIR tradename from Glaxo-Wellcome Research Triangle, NC 27709; abacavir (1592U89) disclosed in WO96/30025 and available under the ZIAGEN tradename from Glaxo-Wellcome Research Triangle, NC 27709; acyclovir dipivoxil [bis(POM)-PMEA] available under the PREVON tradename from Glaxo Sciences, Foster City, CA 94404; lobucavir (BMS-180194), a nucleoside reverse transcriptase inhibitor disclosed in EP-0358154 and EP-0736533 and under development by Bristol-Myers Squibb, Princeton, NJ 08543; BCH-10652, a reverse transcriptase inhibitor (in the form of a racemic mixture of BCH-10618 and BCH-10619) under development by Biochem Pharma, Laval, Quebec H7V, 4A7, Canada; emtricitabine ((-)-FTC) licensed from Emory University under Emory Univ. U.S. Patent No. 5,814,639 and under development by Triangle Pharmaceuticals, Durham, NC 27707; beta-L-FD4(also called beta-L-D4C and named beta-L-2', 3'-dideoxy-5-fluorocytidine) licensed by Yale University to Vion Pharmaceuticals, New Haven CT 06511; and DAPD, the purine nucleoside, (-)-beta-D-2,6,-diaminopurine dioxolane disclosed in EP 0656778 and licensed by Emory University and the University of Georgia to Triangle Pharmaceuticals, Durham, NC 27707; and Iodenosine (FddA), 9-(2,3-dideoxy-2-fluoro-b-D-threo-pentofuranosyl)adenine, a acid stable purine-based reverse transcriptase inhibitor discovered by the NIH and under development by U.S. Bioscience Inc., West Conshohocken, PA, 19428.

The term "non-nucleoside reverse transcriptase inhibitors" ("NNRTI"s) as used herein means non-nucleosides that inhibit the activity of HIV-1 reverse transcriptase.

Typical suitable non-nucleoside reverse transcriptase inhibitors include nevirapine (BI-RG-587) available under the VIRAMUNE tradename from Boehringer Ingelheim, the manufacturer for Roxane Laboratories, Columbus, OH

27

- 43216; delaviradine (BHAP, U-90152) available under the RESCRIPTOR tradename from Pharmacia & Upjohn Co., Bridgewater NJ 08807; efavirenz (DMP-266) a benzoxazin-2-one disclosed in WO94/03440 and available under the SUSTIVA tradename from DuPont Pharmaceutical Co., Wilmington, DE 19880-0723; DuPont 961 and DuPont 083 also under development by DuPont Pharmaceutical Co., Wilmington, DE 19880-0723; PNU-142721, a furopyridine-thiopyrimidine under development by Pharmacia and Upjohn, Bridgewater NJ 08807; capravirine(formerly AG-1549 or Shionogi # S-1153); 5-(3,5-dichlorophenyl)-thio-4-isopropyl-1-(4-pyridyl)methyl-H-imidazo[2-y]methyl carbonate disclosed in WO 96/10019 and under clinical development by Agouron Pharmaceuticals, Inc., LaJolla CA 92037-1020; MKC-442 1-(ethoxymethyl)-5-(1-methylethyl)-6-(phenylmethyl)-(2,4(1H,3H)-pyrimidinedione discovered by Mitsubishi Chemical Co. and under development by Triangle Pharmaceuticals, Durham, NC 27707; and (+)-calanolide A (NSC-675451) and B coumarin derivatives disclosed in NIH U.S. Patent No. 5,489,697, licensed to Med Chem Research, which is co-developing (+) calanolide A with Vira-Invest as an orally administrable product.

- The term "protease inhibitor" ("PI") as used herein means inhibitors of the HIV-1 protease, an enzyme required for the proteolytic cleavage of viral polypeptide precursors (e.g., viral GAG and GAG Pol polyproteins), into the individual functional proteins found in infectious HIV-1. HIV protease inhibitors include compounds having a peptidomimetic structure, high molecular weight (7600 daltons) and substantial peptide character, e.g., CRUXIVAN(available from Merck) as well as nonpeptide protease inhibitors e.g., VIRACEPT (available from Agouron).

- Typical suitable protease inhibitors include saquinavir (Ro 31-8959) available in hard gel capsules under the INVIRASE tradename and as soft gel capsules under the FORTOUASE tradename from Roche Pharmaceuticals, Nutley, NJ 07110-1199; ritonavir (ABT-538) available under the

28

- NORVIR tradename from Abbott Laboratories, Abbott Park, IL 60064; indinavir (MK-639) available under the CRUXIVAN tradename from Merck & Co., Inc., West Point, PA 19486-0004; MK-994, under development by Merck & Co., Inc., West Point, PA 19486-0004; neviravir (AG-1343) available under the VIRACEPT tradename from Agouron Pharmaceuticals, Inc., LaJolla, CA 92037-1020; Ag 1776 under development by Agouron Pharmaceuticals, Inc., LaJolla, CA 92037-1020; amprenavir (141W94), a non-peptide protease inhibitor, tradename AGENERASE, under development by Vertex Pharmaceuticals, Inc., Cambridge, MA 02139-4211 and available from Glaxo-Wellcome, Research Triangle, NC under an expanded access program; zalcitabine (BMS-234475) available from Bristol-Myers Squibb, Princeton, NJ 08543 (originally discovered by Novartis, Basel, Switzerland (CGP-61755); DMP-450, a cyclic urea discovered by Dupont and under development by Triangle Pharmaceuticals; BMS-2322632, PD 178390 under development by Pake-Davis, Morris Plains, NJ 07050; an azapeptide under development by Bristol-Myers Squibb, Princeton, NJ 08543 as a 2nd-generation HIV-1 PI; and ABT-378 under development by Abbott, Abbott Park, IL 60064; and AG-1549 an orally active imidazole carbamate discovered by Shionogi (Shionogi #S-1153) and under development by Agouron Pharmaceuticals, Inc., LaJolla CA 92037-1020.

- The term "anti-HIV-1 therapy" as used herein means any anti-HIV-1 drug found useful for treating HIV-1 infections in man alone, or as part of multidrug combination therapies, especially the triple and quadruple combination therapies called HAART.

- Typical suitable anti-HIV-1 therapies include, but are not limited to multidrug combination therapies such as (i) at least three anti-HIV-1 drugs selected from two NRTIs, one PI, a second PI, and one NNRTI; and (ii) at least two anti-HIV-1 drugs selected from, NNRTIs and PIs; see Tables I, II and III, hereinafter.

29

Typical suitable HAART - multidrug combination therapies- include

- (a) triple combination therapies such as two NRTIs and one PI ; or (b) two NRTIs and one NNRTI ; and (c) quadruple combination therapies such as two NRTIs , one PI and a second PI or one NNRTI. In treatment-naïve patients, it is preferred to start anti-HIV-1 treatment with the triple combination therapy; the use of two NRTIs and one PI is preferred unless there is intolerance to PIs. Drug compliance is essential. The CD4⁺ and HIV-1-RNA plasma levels should be monitored every 3-6 months. Should viral load plateau, a fourth drug e.g., one PI or one NNRTI could be added. See the Table A hereinbelow.

Table A

ANTI-HIV-1 MULTI DRUG COMBINATION THERAPIES

A. Triple Combination Therapies

1. Two NRTIs¹ + one PI²
2. Two NRTIs¹ + one NNRTI³

B. Quadruple Combination Therapies⁴

- 20 Two NRTIs + one PI + a second PI or one NNRTI

C. ALTERNATIVES,⁵

- 25 Two NTRI¹
- One NTRI² + one PI²
- Two PIs² ± one NTRI¹ or NNRTI³
- One PI² + one NRTI¹ + one NNRTI³

30

FOOTNOTES TO TABLE A

1. One of the following: zidovudine + lamivudine; zidovudine + didanosine; stavudine + lamivudine; stavudine + didanosine; zidovudine + zalcitabine; See also Table I
- 5 2. Indinavir, nelfinavir, ritonavir or saquinavir soft gel capsules. Ritonavir is used less frequently because of troublesome adverse effects. The old formulation of saquinavir was used least often because of its poor bioavailability and limited effectiveness, but the new saquinavir formulation should be more effective. See also Table III.
- 10 3. Nevirapine or delavirdine. See also Table II
4. See A-M. Vandamme et al Antiviral Chemistry + Chemotherapy 9:187 at p 193-197 and Figures 1 + 2.
5. Alternative regimens are for patients unable to take a recommended regimen because of compliance problems or toxicity, and for those who fail or relapse on a recommended regimen. Double nucleoside combinations may lead to HIV- resistance and clinical failure in many patients.
6. Most data obtained with saquinavir and ritonavir (each 400 mg bid). See also Table III
7. Zidovudine, stavudine or didanosine. See also Table I
- 20 Other anti-HIV-1 drugs useful for administration in association with pegylated interferon alfa include hydroxyurea, ribavirin, IL-2 and IL-12, and Yissum Project No. 11607. These above-listed anti-HIV-1 drugs may also be administered in association with pegylated interferon alfa in association with any anti-HIV-1 drug therapy, especially the triple and quadruple drug combinations called HAART.
- 25 Hydroxyurea (Droxia) is a ribonucleoside triphosphate reductase inhibitor, the enzyme involved in the activation of T-cells. Hydroxyurea discovered at the NCI is under development by Bristol-Myers Squibb. In preclinical studies, it was
- 30

31

shown to have a synergistic effect on the activity of didanosine and has been studied with stavudine.

- 5 Yissum Project No. 11607, a synthetic protein based on the HIV -1 Vif protein under preclinical development by Yissum Research Development Co., Jerusalem 91042, Israel.

- 10 IL-2 is disclosed in Ajinomoto EP-0142268, Takeda EP-0176299, and Chiron U. S. Patent Nos. RE 33653, 4530787, 4569790, 4604377, 4748234, 4752585, and 4949314 is available under the PROLEUKIN(aldesleukin) tradename from Chiron Corp., Emeryville, CA 94608-2997 as a lyophilized powder for IV infusion or sc administration upon reconstitution and dilution with water, doses of about 1 to about 20 million IU/day, sc is preferred; a dose of about 15 million IU/day, sc is more preferred.

- 15 IL-12 is disclosed in WO96/25171 and is available from Roche Pharmaceuticals, Nutley, NJ 07110-1199 and American Home Products, Madison, NJ 07940; a dose of about 0.5 microgram/kg/day to about 10 microgram/kg/day, sc.

- 20 Pentafuside (DP-178, T-20) a 36-amino acid synthetic peptide disclosed in U.S. Patent No.5,464,933 licensed from Duke University to Trimeris which is developing pentafuside in collaboration with Duke University; pentafuside acts by inhibiting fusion of HIV-1 to target membranes. Pentafuside (3-100 mg /day) is given as a continuous sc infusion or injection together with efavirenz and 2 PIs to HIV-1 positive patients refractory to a triple combination therapy; use of 100 mg/day is preferred. A second generation fusion inhibitor T-1249 (39aa) is under development by Trimeris. Other inhibitors under development include CXCR4, AMO3100, and INTERGRASE by Merck & Co.

30

32

- 5 Ribavirin, 1- β -D-ribofuranosyl-1H-1,2,4-triazole-3-carboxamide, available from ICN Pharmaceuticals, Inc., Costa Mesa, California, is described in the Merck Index, compound No. 8199, Eleventh Edition. Its manufacture and formulation is described in U.S. Patent No. 4,211,771.

- 10 The pegylated interferon α 1a, PEG₁₀₀₀-IFN- α 1a2b (available from Schering-Plough Research Institute, Kenilworth, NJ) increased the LD_{50} anti HIV-1 activity of ribavirin. The combination of PEG₁₀₀₀-IFN- α 1a2b and ribavirin inhibited HIV replication in vitro using phytohemagglutinin ("PHA" - P) - activated peripheral blood mononuclear cells ("PBMCs") at doses corresponding to plasmatic concentrations observed in animals and man. Healthy PBMCs were separated from a buffy-coat of one HIV-seronegative blood donor by Ficoll-Hypaque density gradient centrifugation. PBMCs were activated by 1 μ g/ml phytohemagglutinin (PHA-P) for two days in cell culture medium A: RPMI 1640 supplemented with 10% heat-inactivated (+56°C, 45 min.) fetal calf plasma (FCS), 2 mM L-glutamine and a tri-antibiotic mixture (penicillin, streptomycin, neomycin; PSN). After these two days, cells were washed and cultured at one million cell per milliliter in cell culture medium B: cell culture medium A supplemented with 20 IU/ml recombinant human interleukin-2. Cells were maintained at +37°C in a 5% CO₂ air humidified atmosphere. Experiments were repeated twice with cells of other blood donors. In total, three independent experiments were performed.

- 25 PBMCs were infected with 1,000 50% Tissue Culture Infectious Doses (TCID₅₀) of the reference HIV-1-LAI strain [F. Barre-Sinoussi, Science, 1983, 220, 868-871]. This strain has been amplified using PHA-P-activated umbilical blood mononuclear cells (UBMC). Viral stock has been then titrated on PHA-P activated PBMC by end-point dilution. TCID₅₀ was then calculated using Karber's formula [Arch. Exp. Path. Pharmac., 1931, 162, 126-139].

30

33

PEG₁₂₀₀₀-IFN- α 2b and ribavirin, alone and in combination, and AZT used as a control, were administered 24 hours before HIV-1 infection and maintained all along the culture. Three doses of PEG₁₂₀₀₀-IFN- α 2b and ribavirin were used.

5 200,000 PHA-P-activated PBMCs were added to each well of 96-well microplates. Cells were 24 hour-pretreated prior to infection with the reference HIV-1-LAI strain. Twice a week, cell supernatants were collected, and drugs and medium were renewed. At day 7, RT activity were determined in cell supernatants, and potential cytotoxic effects of drugs and drug combinations were evaluated by microscopic observation.

10 Viral replication was measured by determining reverse transcriptase ("RT") activity in cell supernatants using Retro-Sys® kit, according to manufacturer's recommendations (Imnovagen, Lund, Sweden).

15 Effective doses were calculated using cumulative RT activities with Chou J. and TC: microcomputer software.

20 The combined effects were analyzed using either the combination index (CI) [Chou & Talalay, 1984] with J and TC Chou microcomputer software, or the fractionary inhibitory concentration (FIC) index [Antimicrob. Agents. Chemother., 1987, 31, 1613-1617]. When the CI or FIC index is equal to 1, the combination is additive. When it is below 1.0, the combination is synergistic, and when it is above 1.0, the combination is judged as antagonistic.

25 PEG₁₂₀₀₀-IFN- α 1a2b as well as the combination of PEG₁₂₀₀₀-IFN- α 1a2b and ribavirin inhibited the HIV replication at doses corresponding to plasmatic concentrations measured in mice and HIV-1 infected patients [BE. Gilbert, et al. Antimicrob. Agents Chemother., 1988, 32, 117-121; E. Connor et al., Antimicrob. Agents Chemother., 1993, 37, 537-539].

34

These above-listed anti-HIV-1 drugs may also be administered in association with pegylated interferon α 1a in association with any anti-HIV-1 drug therapy, especially the triple and quadruple drug combinations called HAART.

5 The term "interferon- α 1a" as used herein means the family of highly homologous species-specific proteins that inhibit viral replication and cellular proliferation and modulate immune response. Typical suitable interferon- α 1as include, but are not limited to, recombinant interferon α 1a-2b, such as Intron-A interferon available from Schering Corporation, Kenilworth, N.J., recombinant

10 interferon α 1a-2a such as Roferon interferon available from Hoffmann-La Roche, Nutley, N.J., recombinant interferon α 1a-2C such as Berofer α 1a 2 interferon available from Boehringer Ingelheim Pharmaceutical, Inc., Ridgefield, CT.,

15 interferon α 1a-n1, a purified blend of natural α 1a interferons such as Sumitron available from Sumitomo, Japan or as Wellferon interferon α 1a-n1 (INS) available from the Glaxo-Wellcome Ltd., London, Great Britain, or a consensus

20 α 1a interferon such as those described in U.S. Patent Nos. 4,897,471 and 4,695,623 (especially Examples 7, 8 or 9 thereof) and the specific product available from Amgen, Inc., Newbury Park, CA, or interferon α 1a-n3 a mixture of natural α 1a interferons made by Interferon Sciences and available from the Purdue Frederick Co., Norwalk, CT., under the Alferon Tradename. The use of interferon α 1a-2a or α 1a 2b is preferred. Since interferon α 1a 2b, among all interferons, has the broadest approval throughout the world for treating chronic hepatitis C infection, it is most preferred. The manufacture of interferon α 1a 2b is described in U.S. Patent No. 4,530,901.

25 The term "pegylated interferon α 1a" as used herein means polyethylene glycol modified conjugates of interferon α 1a, preferably interferon α 1a-2a and -2b. The preferred polyethylene-glycol-interferon α 1a-2b conjugate is PEG₁₂₀₀₀-interferon α 1a 2b. The phrases "12,000 molecular weight polyethylene glycol conjugated interferon α 1a" and "PEG₁₂₀₀₀-IFN α 1a" as used herein mean

30

35

conjugates such as are prepared according to the methods of International Application No. WO 95/13090 and containing urethane linkages between the interferon α -2a or -2b amino groups and polyethylene glycol having an average molecular weight of 12000.

5

The preferred PEG₁₂₀₀₀-interferon α -2b is prepared by attaching a PEG polymer to the epsilon amino group of a lysine residue in the IFN α -2b molecule. A single PEG₁₂₀₀₀ molecule is conjugated to free amino groups on an IFN α -2b molecule via a urethane linkage. This conjugate is characterized by the molecular weight of PEG₁₂₀₀₀ attached. The PEG12000-IFN α -2b conjugate is formulated as a lyophilized powder for injection. The objective of conjugation of IFN α with PEG is to improve the delivery of the protein by significantly prolonging its plasma half-life, and thereby provide protracted activity of IFN α .

15

Other interferon α conjugates can be prepared by coupling an interferon α to a water-soluble polymer. A non-limiting list of such polymers include other polyalkylene oxide homopolymers such as polypropylene glycols, polyoxyethylenated polyols, copolymers thereof and block copolymers thereof. As an alternative to polyalkylene oxide-based polymers, effectively non-antigenic materials such as dextran, polyvinylpyrrolidones, polyacrylamides, polyvinyl alcohols, carbohydrate-based polymers and the like can be used. Such interferon α -polymer conjugates are described in U.S. Patent No. 4,766,106, U.S. Patent No. 4,917,888, European Patent Application No. 0 236 987, European Patent Application Nos. 0510 356, 0 593 668 and 0 809 996 (pegylated interferon α -2a) and International Publication No. WO 95/13090.

25

Pharmaceutical composition of pegylated interferon α -suitable for parenteral administration may be formulated with a suitable buffer, e.g., Tris-HCl, acetate or phosphate such as dibasic sodium phosphate/monobasic sodium phosphate buffer, and pharmaceutically acceptable excipients (e.g., sucrose), carriers (e.g. human plasma albumin), toxicity agents (e.g. NaCl), preservatives

30

36

(e.g. thimerosol, cresol or benylalcohol), and surfactants (e.g. tween or polysorbates) in sterile water for injection. The pegylated interferon α -may be stored as lyophilized powders under a refrigeration at 2°-8°C. The reconstituted aqueous solutions are stable when stored between 2° and 8°C and used within 24 hours of reconstitution. See for example U.S. Patent Nos. 4,492,537, 5,762,923 and 5,766,582. The reconstituted aqueous solutions may also be stored in prefilled, multi-dose syringes such as those useful for delivery of drugs such as insulin. Typical suitable syringes include systems comprising a prefilled vial attached to a pen-type syringe such as the NOVOLET Novo Pen available from Novo Nordisk, as well as prefilled, pen-type syringes which allow easy self-injection by the user. Other syringe systems include a pen-type syringe comprising a glass cartridge containing a diluent and lyophilized pegylated interferon α powder in a separate compartment.

5

10

Certain CCR5 antagonist compounds represented by the structural formula I or II or III or IV may exist in different isomeric (e.g., enantiomers, diastereoisomers and atropisomers) forms. The invention contemplates all such isomers both in pure form and in admixture, including racemic mixtures.

15

20

Certain compounds of formulas I or II or III or IV will be acidic in nature, e.g. those compounds which possess a carboxyl or phenolic hydroxyl group. These compounds may form pharmaceutically acceptable salts. Examples of such salts may include sodium, potassium, calcium, aluminum, gold and silver salts. Also contemplated are salts formed with pharmaceutically acceptable amines such as ammonia, alkyl amines, hydroxyalkylamines, N-methylglucamine and the like.

25

30

Certain basic compounds of formulas I or or III or IV also form pharmaceutically acceptable salts, e.g., acid addition salts. For example, the pyrido-nitrogen atoms may form salts with strong acid, while compounds having basic substituents such as amino groups also form salts with weaker acids. Examples of suitable acids for salt formation are hydrochloric, sulfuric, phosphoric, acetic, citric, oxalic, malonic, salicylic, malic, fumaric, succinic,

37

ascorbic, maleic, methanesulfonic and other mineral and carboxylic acids well known to those in the art. The salts are prepared by contacting the free base form with a sufficient amount of the desired acid to produce a salt in the conventional manner. The free base forms may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous NaOH, potassium carbonate, ammonia and sodium bicarbonate. The free base forms differ from their respective salt forms somewhat in certain physical properties, such as solubility in polar solvents, but the acid and base salts are otherwise equivalent to their respective free base forms for purposes of the invention.

All such acid and base salts are intended to be pharmaceutically acceptable salts within the scope of the invention and all acid and base salts are considered equivalent to the free forms of the corresponding compounds of formulas I or II for purposes of the invention.

Compounds of the formula I or II or III or IV are the invention of other inventive entities. Compounds of the formula I or II are disclosed, together with methods of making them, in commonly-owned U.S. Patent Application (Attorney's Docket # IN01031) and compounds of the formula or III or IV in commonly-owned U.S. Patent Application SN(Attorney's Docket # IN01032), each of which were filed on the same date as this application, both of which are hereby incorporated by reference

Compounds of formulas I or II or III or IV can be also made by the procedures known in the art, for example by the procedures described in the following reaction schemes, by the methods described in the examples below, and by using the methods described in WO96/26196, WO98/05292, WO98/10425 and WO98/06697.

For preparing pharmaceutical compositions from the compounds of formula I or II or III or IV, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets and suppositories. The powders and tablets may be comprised of from about 5 to about 95 percent active ingredient. Suitable solid carriers are known in the art, e.g., magnesium carbonate, magnesium stearate, talc, sugar or lactose. Tablets, powders, cachets and capsules can be used as solid dosage forms suitable for oral administration. Examples of pharmaceutically

38

acceptable carriers and methods of manufacture for various compositions may be found in A. Gennaro (ed.), Remington's Pharmaceutical Sciences, 18th Edition, (1990), Mack Publishing Co., Easton, Pennsylvania.

Liquid form preparations include solutions, suspensions and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injection or addition of sweeteners and opacifiers for oral solutions, suspensions and emulsions. Liquid form preparations may also include solutions for intranasal administration.

Aerosol preparations suitable for inhalation may include solutions and solids in powder form, which may be in combination with a pharmaceutically acceptable carrier, such as an inert compressed gas, e.g., nitrogen.

Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions and emulsions.

The compounds of formula I or II or III or IV may also be deliverable transdermally. The transdermal compositions can take the form of creams, lotions, aerosols and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type as are conventional in the art for this purpose.

Preferably the compounds of formula I or II or III or IV is administered orally.

Preferably, the pharmaceutical preparation is in a unit dosage form. In such form, the preparation is subdivided into suitably sized unit doses containing appropriate quantities of the active component, e.g., an effective amount to achieve the desired purpose.

The quantity of active compound of formula I or II or III or IV in a unit dose of preparation may be varied or adjusted from about 10 mg to about 500 mg, preferably from about 25 mg to about 300 mg, more preferably from about 50 mg to about 250 mg, and most preferably from about 55 mg to about 200 mg, according to the particular application.

The actual dosage employed may be varied depending upon the requirements of the patient and the severity of the condition being treated. Determination of the proper dosage regimen for a particular situation is within the skill of the art. For convenience, the total daily dosage may be divided and administered in portions during the day as required.

39

The amount and frequency of administration of the compounds of formula I or II or III or IV and/or the pharmaceutically acceptable salts thereof will be regulated according to the judgment of the attending clinician considering such factors as age, condition and size of the patient as well as severity of the symptoms being treated. A typical recommended daily dosage regimen for oral administration of the compounds of formula I or II or III or IV can range from about 100 mg/day to about 300 mg/day, preferably 150 mg/day to 250 mg/day, more preferably about 200 mg/day, in two to four divided doses.

10 A person suffering from chronic hepatitis C infection may exhibit one or more of the following signs or symptoms:

(a) elevated ALT,

15 (b) positive test for anti-HCV antibodies,

(c) presence of HCV as demonstrated by a positive test for the presence of HCV-RNA in the serum,

20 (d) clinical stigmata of chronic liver disease,

(e) hepatocellular damage.

25 In a preferred aspect of the present invention, a therapeutically effective amount of the combination therapy of pegylated interferon- α and a CCR5 antagonist compound represented by structural formula I or II or III or IV is administered in association with a therapeutically effective amount of ribavirin and anti-retroviral therapy, e.g., HAART, to the patient having HIV-1 infection and exhibiting one or more of the above signs or symptoms in the first and second treatment time periods in amounts sufficient to eliminate or at least alleviate one or more of the signs or symptoms, and to lower the HCV-RNA plasma levels by at least a power of ten, and preferably to eradicate detectable HCV-RNA at least

40

by the end of the second treatment time period and to maintain no detectable HCV-RNA for at least 24 weeks after the end of the second treatment time period. The sum of the first and second treatment time periods is about 40-50 weeks, and preferably is 48 weeks. Administration of the ribavirin may be discontinued after the end of the second time period depending upon the judgment of the attending clinician.

10 The term "no detectable HCV-RNA" in the context of the present invention means that there are fewer than 100 copies of HCV-RNA per ml of plasma of the patient as measured by quantitative, multi-cycle reverse transcriptase PCR methodology. HCV-RNA is preferably measured in the present invention by research-based RT-PCR methodology well known to the skilled clinician. This methodology is referred to herein as HCV-RNA/qPCR. The lower limit of detection of HCV-RNA is 100 copies/mL. Serum HCV-RNA/qPCR testing and HCV genotype testing will be performed by a central laboratory. See also J. G. McHutchinson et al. (N. Engl. J. Med., 1998, 339:1485-1492), and G. L. Davis et al. (N. Engl. J. Med. 339:1493-1499).

20 In a preferred embodiment of the present invention, those patients co-infected with HIV-1 and HCV infections are treated with a combination therapy of pegylated interferon α and a CCR5 antagonist compound represented by structural formula I or II or III or IV in association with ribavirin and a HAART combination considered appropriate by the attending clinician and the patient; use of the interferon α -2b-ribavirin combination therapy sold by Schering Corp. under the REBETRON trademark is preferred. See also J. G. McHutchinson et al. (N. Engl. J. Med., 1998, 339:1485-1492), and G. L. Davis et al. (N. Engl. J. Med. 339:1493-1499). Ribavirin, 1- β -D-ribofuranosyl-1H-1,2,4-triazole-3-carboxamide, available from ICN Pharmaceuticals, Inc., Costa Mesa, California, is described in the Merck Index, compound No. 8199, Eleventh Edition. Its manufacture and formulation is described in U.S. Patent No. 4,211,771.

41

For the pediatric patient co-infected with the H IV-1 and HCV infections, a suitable HAAART includes a NRTI+ a PI, e.g., Nelfinavir +a NNRTI, e.g., Efavirenz in combination with the dosages and dosage regimens for pegylated interferon alfa and ribavirin listed herein above. See also Tables I-IV herein below. A human growth hormone such as the polypeptide hormone, somatropin, of recombinant rDNA origin, available under the HUMATROPE tradename from Eli Lilly & Co., Indianapolis, IN 46285, may be administered to these pediatric patients in the dosage and administration schedule listed in the product information sheet in consultation with the attending clinician to reduce retardation of growth associated with pegylated interferon alfa treatment.

HAAART is administered to the patient in association with pegylated interferon-alfa, that is, the pegylated interferon-alfa dose may be administered before, after or during the same period of time that the patient receives doses of HAAART. A human growth hormone such as the polypeptide hormone, somatropin, of recombinant rDNA origin, available under the HUMATROPE tradename from Eli Lilly & Co., Indianapolis, IN 46285, may also be administered in association with HAAART and pegylated interferon alfa- to the pediatric patient having HIV-1 infection in the dosage and administration schedule listed in the product information sheet in consultation with the attending clinician.

In a preferred embodiment of the present invention, pegylated interferon alfa is administered to HIV-1 infected patients prior to initiation of a CCR5 antagonist compound represented by structural formula I or II or III or IV and of HAAART, and preferably about two to about four weeks prior to initiation of HAAART. In another preferred embodiment of the present invention, administration of pegylated interferon alfa is initiated concurrently, i.e., on the same day with the administration of a CCR5 antagonist compound represented by structural formula I or II or III or IV and HAAART. In another preferred embodiment of the present invention the pegylated interferon-alfa is administered

42

after the HIV-1 infected patient has initiated use of a CCR5 antagonist compound represented by structural formula I or II or III or IV and HAAART.

The goal of the HIV-1 therapy of the present invention is to reduce the HIV-1-RNA viral load below the detectable limit. The "detectable limit of HIV-1-RNA" in the context of the present invention means that there are fewer than about 200 to fewer than about 50 copies of HIV-1-RNA per ml of plasma of the patient as measured by quantitative, multi-cycle reverse transcriptase PCR methodology.

HIV-1-RNA is preferably measured in the present invention by the methodology of Amplicor -1 Monitor 1.5 (available from Roche Diagnostics) or of Nuclisens HIV-1 QT -1. This methodology is described by Schooley, RT, Antiviral Therapy(1997), 2 (Suppl. 4):59-70.

The doses and dosage regimen of the NRTIs, NNRTIs, PI, pentastide, IL-2, IL-12, a CCR5 antagonist compound represented by structural formula I or II or III or IV and pegylated interferon alfa will be determined by attending clinician in view of the approved doses and dosage regimen in the package insert or as set forth in the protocol taking into consideration the age, sex and condition of the patient and the severity of the HIV-1 and HCV infections. For the pediatric patient infected with the H IV-1, or co-infected with the H IV-1 and HCV infections a suitable HAAART includes a NRTI+ a PI, e.g., Nelfinavir +a NNRTI, e.g., Efavirenz in combination with the dosages and dosage regimens for pegylated interferon alfa and ribavirin listed herein above. See also Tables I-IV hereinafter for dosages and dosage regimens.

The following clinical protocol may be used to administer the anti-HIV-1 therapy of the present invention. Many modifications of this clinical protocol will be obvious to the skilled clinician, and the following Study Design should not be interpreted as limiting the scope of the method of this invention which is only limited by the claims listed hereinafter. See for example J. G. McInturk et al.

(N. Engl. J. Med., 1998, 339:1485-1492), and G. L. Davis et al. (N. Engl. J. Med. 339:1493-1499).

The study population should include male and female patients

- 5 diagnosed with HIV-1 infection who are either treatment naive or treatment-experienced and should be included if they meet the following inclusion and exclusion criteria:

10 **Subject Inclusion Criteria:**

- Subjects diagnosed with HIV-1 infection who are either treatment naive or treatment-experienced.
- HIV-RNA by Amplicor test, Version 1.5 of greater than 500 copies/mL.
- 15 • CD₄ count greater than 100 copies/mL, preferably greater than 200 cells/mL.
- Subjects in good physical health with clinically acceptable safety laboratory test results and ECG.
- The following laboratory parameters must be met:
- Platelet count •100,00/mL
- 20 • Hemoglobin •9 gm/dL (90 gm/L)
- Absolute neutrophil count •1500/ μ L
- Creatinine <1.5 times the upper limit of normal
- SGOT/SGPT \leq 5 x upper limit of normal
- Bilirubin \leq 2.5 x upper limit of normal
- 25 • A negative urine pregnancy test (females only)

Subjects must be willing and able to give written informed consent and be able to adhere to the schedule set forth in the protocol.

Subject Exclusion Criteria

- Females who are breast-feeding or pregnant or who are not using adequate birth control.


- Subject with allergy to E. coli proteins
- Subjects with a significant past medical/psychiatric history, specifically depression or dementia.

- 5 In a preferred embodiment of the present invention, the subjects will be randomized to receive pegylated interferon alfa 2b, i.e., PEG₁₂₀₀-interferon alfa 2b at doses between 0.5 and 4.5 micrograms per kilogram e.g. at doses of 0.5, 1.0, 1.5, 3.0 or 4.5 micrograms per kilogram by subcutaneous injection once a week.
- 10 The amount of ribavirin administered concurrently with the pegylated interferon-alfa is from about 400 to about 1600 mg per day, preferably about 600 to about 1200 mg/day or about 800 to about 1200 mg day and most preferably about 1000 to about 1200 mg/kg a day.

- 15 The quantity of CCR5 antagonist compound of formula I or II or III or IV in a unit dose or formulation may be varied or adjusted from about 10 mg to about 500 mg, preferably from about 25 mg to about 300 mg, more preferably from about 50 mg to about 250 mg, and most preferably from about 55 mg to about 200 mg, according to the particular application.

- 20 HAART may also be initiated before or concurrently with the administration of the pegylated interferon alfa 2b, i.e., PEG₁₂₀₀-interferon alfa 2b, a CCR5 antagonist compound of formula I or II or III or IV and ribavirin.


- 25 CCR5 antagonist compounds of the following structures are representative of formulas I and II useful in the present invention:



5 wherein R⁶, X and R² are as defined in the following table:

R ⁶	X	R ²
Br		
Br		
Br		
Br		
Br		
Br		
F ₃ C-		

CCR5 antagonist compounds of the following structures are representative of formulas III and IV useful in the present invention:



5 wherein R, R³, R⁶ and R² are as defined in the following table:

R	R ³	R ⁶	R ²
		-CH ₃	
		-CH ₃	
	H	-CH ₃	
		H	
		-CH ₃	
		-CH ₃	

The actual dosages employed may be varied depending upon the requirements of the patient and the severity of the condition being treated. Determination of the proper dosage regimen for a particular situation is within the skill of the art. For convenience, the total daily dosage may be divided and administered in portions during the day as required.

Overall Design and Plan of the Study:
The primary efficacy objective will be lowering of the HIV-1-RNA plasma levels by a factor of 10 or greater.

Plasma HIV-1-RNA/qPCR testing will be performed by a central laboratory. A positive HIV-1-RNA assay result will be required at Baseline, only patients positive for HIV-1-RNA will be eligible to participate.

Table I
NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS (NRTI) DOSAGE & DOSAGE REGIMEN

	<u>NRTI (Trade name, Marketer)</u>	<u>Usual adult dosage</u>
10	Zidovudine, AZT (Retrovir - Glaxo Wellcome)*	200 mg PO tid or 300 mg PO bid
	Stavudine (Zerit - Bristol-Myers Squibb)*	40 mg PO bid ¹
15	Didanosine (Videx - Bristol-Myers Squibb)*	200 mg PO bid ²
	Lamivudine (Epivir - Glaxo Wellcome)*	150 mg PO bid ³
20	Zalcitabine (Hivid - Roche)	0.75 mg PO tid
	Zidovudine plus lamivudine (Combivir - Glaxo Wellcome)	1 tablet PO bid ⁴
25	Abacavir (Ziagen-Glaxo-Wellcome)	200 or 400 mg PO tid
	Adefovir dipivoxil (Prevon-Gilead Sciences)	125 or 200 mg PO qd ⁵
30	Lobucavir (BMS-180194-BMS)	200 mg PO bid ⁶
	BCH-10652 (Biochem Pharme)	200 mg PO, qid 400 mg PO, qid ⁷
	Emtricitabine ((-)-FTC-Triangle Pharmaceuticals)	200 mg PO qd ⁸
35	Beta-L-FD4 (B-L-D4C-Vion Pharmaceutical)	0.2-25 mg/kg/day ⁹
	DAPD (Triangle Pharmaceuticals)	¹⁰
40	Lodenosine (Fdda-U.S. Bioscience)	1.6-3.2 mg/Kg PO bid ¹¹

Footnotes Table I

¹ Available in a liquid formulation.
² For patients less than 60 kg, 30 mg PO bid.

³ With tablets, for patients < 60 kg, 125 mg PO bid, > 60 kg, 200 mg PO bid. With powder, dosage varies from 167 mg (< 60 kg) to 250 mg PO (< 60 kg) bid. Doses should be taken at least 30 minutes before meals or at least two hours afterward.

Footnotes Table I

⁴ For patients less than 50 kg, 2 mg/kg PO bid.
⁵ Each tablet contains 300 mg of zidovudine and 150 mg of lamivudine.
⁶ Available under an expanded access program - a NIH-sponsored Phase III Trial
⁷ Phase II.
⁸ Phase I/II; see PharmaProjects, sections J5A & J5Z.
⁹ Phase I/II; see PharmaProjects, sections J5A & J5Z.
¹⁰ Preclinical; active in duck HBV model; see PharmaProjects, sections J5A & J5Z.
¹¹ Preclinical; active po and IV; DAPD is a prodrug of another dioxolene purine, DXG. See PharmaProjects, sections J5A & J5Z.
Phase II, Fdda has potential for once-a-day dosage.

TABLE II

NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS (NNRTI) Dosage and Dosage Regimen

	Usual adult dosage and Dosage Regimen
5 <u>NNRTI (Tradename, Marketer)</u>	
10 Nevirapine (Viramune - Roxane)	200 mg PO bid ¹
Delavirdine (Rescriptor - Pharmacia & Upjohn)	400 mg PO tid
Efavirenz (Sustiva, Dupont)	200 mg PO qid ²
15 PNU-142721 (Pharmacia + Upjohn)	_____
AG-1549 (Agouvon Pharmaceuticals)	_____
MKC-442 (Triangle Pharmaceuticals)	750 mg PO bid ³
20 (+)-Calanoides A (Med Chem Research)	800 mg PO ⁴
25	
30	

For the first two weeks of treatment with nevirapine, to decrease the risk of rash, patients should take only one 200-mg tablet per day.

Quadruple Therapy of efavirenz with didanosine + AZT + lamivudine.

Phase I/II evaluating dose and concomitant use with other anti-HIV-1 therapies; see Pharmacokinetics, sections J5A & J5Z.

Triple Therapy of (a) MKC-442 with stavudine and either lamivudine or didanosine or (b) MKC-442 with nevirapine (qv) and NRTIs.

Phase I; see Pharmacokinetics, sections J5A & J5Z.

TABLE III

Protease Inhibitor (PI) Dosage and Dosage Regimen

	Dosage + Dosage Regimen
5 <u>PI (Tradename, Marketer)</u>	
10 Saquinavir (Invirase - hard gel capsule - Roche) (Fortovase - soft gel capsule - Roche)	600 mg PO tid ¹ 1100 mg PO tid ¹
10 Ritonavir (Norvir - Abbott)	600 mg PO bid ²
Indinavir (Crivivan - Merck)	800 mg PO qid ³
Nelfinavir (Viracept - Agouron)	750 mg PO tid ⁴
15 Agenerase (Amprenavir, 141W94, Glaxo)	900 mg - 1200 mg PO bid ⁵
Lasinavir (BMS-234475, BMS)	_____
20 DMP-450 (Triangle Pharmaceuticals)	_____
BMS-2322623 (BMS)	_____
25 ABT-378 (Abbott)	60 mg PO bid ⁶
30	
35	
40	

With, or within two hours after, a full meal.

With food. The liquid formulation has an unpleasant taste; the manufacturer suggests taking it with chocolate milk or a liquid nutritional supplement.

With water, one hour before or two hours after a meal. Patients taking indinavir should drink at least 48 ounces (1.5 liter) of water daily.

With food.

Quadruple Combination Therapy of amprenavir with AZT + lamivudine + abacavir.

Phase I/II; see Pharmacokinetics, sections J5A & J5Z.

Phase II; see Pharmacokinetics, sections J5A & J5Z.

Preclinical Studies; Prodrug esters of BMS 2322623 enhance oral absorption; see Pharmacokinetics, sections J5A & J5Z.

Phase I Studies show ABT-378 to be 10X more potent than ritonavir; see Pharmacokinetics sections J5A & J5Z.

51

TABLE IV

Other Anti-HIV-1 Drugs

Drug (Trade Name, Marketer)	Usual Adult Dosage and Dosage Regimen
Hydroxyurea (Droxia, BMS)	1000 mg PO qid ¹
Ribavirin (Rebetol, Schering-Plough)	600mg-1200mg/day, PO
IL-2 (Proleukin, Chiron Corp.)	1-20million IU/day, sc
IL-12 (Roche)	0.5-10 micrograms/kg/day, sc
Yissum Project No. 11607 (Yissum)	_____ ²
Triple Therapy of hydroxyurea with 400 mg qid + 500 mg AZT; see PharmaProjects, section B3C1 Preclinical; see PharmaProjects, sections J5A & J5Z.	

25

52

The following assays can be used to identify a compound as a CCR5 antagonist as well as to determine the CCR5 antagonistic activity of the compounds of formulas I to IV. These assays are disclosed in commonly-owned U.S. Patent Application (Attorney's Docket # INO1031) and commonly-owned U.S. Patent Application SN/Attorney's Docket # INO1032, filed on the same date as this application

CCR5 Membrane Binding Assay:

A high throughput screen utilizing a CCR5 membrane binding assay identifies inhibitors of RANTES binding. This assay utilizes membranes prepared from NIH 3T3 cells expressing the human CCR5 chemokine receptor which have the ability to bind to RANTES, a natural ligand for the receptor. Using a 96-well plate format, membrane preparations are incubated with ¹²⁵I-RANTES in the presence or absence of compound for one hour. Compounds are serially diluted over a wide range of 0.001ug/ml to 1 ug/ml and tested in triplicates. Reaction cocktails are harvested through glass fiber filters, and washed thoroughly. Total counts for replicates are averaged and data reported as the concentration required to inhibit 50 percent of total ¹²⁵I-RANTES binding. Compounds with potent activity in the membrane binding assay are further characterized in second day cell-based HIV-1 entry and replication assays.

HIV-1 Entry Assay:

Replication defective HIV-1 reporter virions are generated by cotransfection of a plasmid encoding the NL4-3 strain of HIV-1 (which has been modified by mutation of the envelope gene and introduction of a luciferase reporter plasmid) along with a plasmid encoding one of several HIV-1 envelope genes as described by Connor et al., *Virology*, 206 (1995), p. 935-944. Following transfection of the two plasmids by calcium phosphate precipitation, the viral supernatants are harvested on day 3 and a functional viral titer determined. These stocks are then used to infect U87 cells stably expressing CD4 and the chemokine receptor CCR5 which have been preincubated with or without test compound. Infections are carried out for 2 hours at 37 °C, the cells washed and media replaced with fresh media containing compound. The cells are incubated for 3 days, lysed and luciferase activity determined. Results are reported as the concentration of compound required to inhibit 50% of the luciferase activity in the control cultures.

HIV-1 Replication Assay:

This assay uses primary peripheral blood mononuclear cells or the stable U87-CCR5 cell line to determine the effect of anti-CCR5 compounds to block infection of primary HIV-1 strains. The primary lymphocytes are purified from normal healthy donors and stimulated *in vitro* with PHA and IL-2 three days prior to infection. Using a 96-well plate format, cells are pretreated with drug for 1 hour at 37 °C and subsequently infected with an M-tropic HIV-1 isolates. Following infection, the cells are washed to remove residual inoculum and cultured in the presence of compound for 4 days. Culture supernatants are harvested and viral replication measured by determination of viral p24 antigen concentration.

10 Calcium Flux Assay:

Cells expressing the HIV coreceptor CCR5 are loaded with calcium sensitive dyes prior to addition of compound or the natural CCR5 ligand. Compounds with agonist properties will induce a calcium flux signal in the cell, while CCR5 antagonists are identified as compounds which do not induce signaling by themselves but are capable of blocking signaling by the natural ligand RANTES.

15 GTP/S Binding Assay:

A GTP/S binding assay measures receptor activation by CCR5 ligands. This assay measures the binding of ³⁵S labeled-GTP to receptor coupled G-proteins that occurs as a result of receptor activation by an appropriate ligand. In this assay, the CCR5 ligand, RANTES, is incubated with membranes from CCR5 expressing cells and binding to the receptor activation (or binding) is determined by assaying for bound ³⁵S label. The assay quantitatively determines if compounds exhibit agonist characteristics by inducing activation of the receptor or alternatively antagonist properties by measuring inhibition of RANTES binding in a competitive or non-competitive fashion.

25 Chemotaxis Assay:

The chemotaxis assay is a functional assay which characterizes the agonist vs. antagonist properties of the test compounds. The assay measures the ability of a non-adherent murine cell line expressing human CCR5 (BaF-550) to migrate across a membrane in response to either test compounds or natural ligands (i.e., RANTES, MIP-1 β). Cells migrate across the permeable membrane towards compounds with agonist activity. Compounds that are antagonists not only fail to induce chemotaxis, but are also capable of inhibiting cell migration in response to known CCR5 ligands.

35


In the assay to determine inhibition of RANTES binding, compounds of the formulas I -IV range in activity from a K_i of about 0.5 to about 1500 nM, with preferred compounds having a range of activity from about 0.5 to about 750 nM, more preferably about 0.5 to 300 nM, and most preferably about 0.5 to 50 nM.

5

We claim:


1. The use of a pegylated interferon- α and a CCR5 antagonist for the preparation of a medicament for the treatment of HIV-1 infections in patients.

2. The use of a pegylated interferon- α and a CCR5 antagonist for the preparation of a medicament for the treatment of HIV-1 infections in patients wherein the CCR5 antagonist is represented by the structural formula I or II or III or IV:






- or a pharmaceutically acceptable salt of I or II or III or IV,

wherein in the CCR5 antagonist compounds represented by structural formula I:



20. X is -C(R¹³)₂-, -C(R¹³)(R¹⁹)-, -C(O)-, -O-, -NH-, -N((C₁-C₆alkyl))-,




- 10 R is R⁶-phenyl, R⁶-pyridyl, R⁶-thiophenyl or R⁶-naphthyl;

R¹ is hydrogen, C₁-C₆ alkyl or C₂-C₆ alkenyl;

R² is R⁷, R⁸, R⁹-phenyl; R⁷, R⁸, R⁹-substituted 6-membered heteroaryl;

R7, R8, R9-substituted 6-membered heteroaryl N-oxide;

R¹⁰, R¹¹-substituted 5-membered heteroaryl; naphthyl; fluorenyl;



R³ is R⁶-phenyl, R⁶-heteroaryl or R⁶-naphthyl;

R⁴ is hydrogen, C₁-C₆ alkyl, fluoro-C₁-C₆ alkyl, cyclopropylmethyl,


$$-\text{CH}_2\text{CH}_2\text{OH}, -\text{CH}_2\text{CH}_2\text{O}-(\text{C}_1-\text{C}_6)\text{alkyl}, -\text{CH}_2\text{C}(\text{O})-\text{O}-(\text{C}_1-\text{C}_6)\text{alkyl},$$
$$-\text{CH}_2\text{C}(\text{O})\text{NH}_2, -\text{CH}_2\text{C}(\text{O})-\text{NH}(\text{C}_1-\text{C}_6\text{alkyl}) \text{ or } -\text{CH}_2\text{C}(\text{O})-\text{N}((\text{C}_1-\text{C}_6\text{alkyl})_2,$$


20 R5 and R1' are independently selected from the group consisting of hydrogen and (C₁-C₆)-alkyl;

R⁶ is 1 to 3 substituents independently selected from the group consisting of hydrogen, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, -CF₃, CF₃O-, CH₃CO)-, -CN,

57

CH₃SO₂-, CF₃SO₂-, R¹⁴-phenyl, R¹⁴-benzyl, CH₃C(=NOCH₃)-,

CH₃C(=NOCH₂CH₃)-, , -NH₂, -NHCOCF₃, -NHCONH(C₁-C₆ alkyl), -NHCO(C₁-C₆ alkyl), -NHCO₂(C₁-C₆ alkyl),

5-membered heterocaryl and , wherein X is -O-, -NH- or -N(CH₃)-;

5 R⁷ and R⁸ are independently selected from the group consisting of (C₁-C₆)alkyl, halogen, -NR²⁰R²¹, -OH, -CF₃, -OCH₃, -O-acyl, and -OCF₃;

R⁹ is R¹, hydrogen, phenyl, -NO₂, -CN, -CH₂F, -CHF₂, -CHO, -CH=NOR²⁰, pyridyl, pyridyl N-oxide, pyrimidinyl, pyrazinyl,

10 -N(R²⁰)CONR²¹R²², -NHCONH(chloro-(C₁-C₆)alkyl), -NHCONH((C₃-C₁₀)cycloalkyl/(C₁-C₆)alkyl), -NHCO(C₁-C₆)alkyl, -NHCOCF₃, -NHSO₂N((C₁-C₆)alkyl)₂, -NHSO₂(C₁-C₆)alkyl, -N(SO₂CF₃)₂, -NHCO₂(C₁-C₆)alkyl, C₃-C₁₀ cycloalkyl, -SR²³, -SOR²³, -SO₂R²³, -SO₂NH(C₁-C₆ alkyl), -OSO₂(C₁-C₆)alkyl, -OSO₂CF₃, hydroxy(C₁-C₆)alkyl, -CON R²⁰R²¹, -CON(CH₂CH₂-O-CH₃)₂, -OC(=O)NH(C₁-C₆)alkyl, -CO₂R²⁰, -Si(CH₃)₃ or -B(OC(CH₃)₂)₂;

15 R¹⁰ is (C₁-C₆)alkyl, -NH₂ or R¹²-phenyl;

R¹² is 1 to 3 substituents independently selected from the group consisting of hydrogen, (C₁-C₆) alkyl, -CF₃, -CO₂R₂₀, -CN, (C₁-C₆)alkoxy and halogen;

R¹³, R¹⁴, R¹⁵ and R¹⁶ are independently selected from the group consisting of hydrogen and (C₁-C₆)alkyl;

20 R¹⁷ and R¹⁸ are independently selected from the group consisting of hydrogen and C₁-C₆ alkyl, or R¹⁷ and R¹⁸ together are a C₂-C₅ alkylene group and with the carbon to which they are attached form a spiro ring of 3 to 6 carbon atoms;


R¹⁹ is R⁶-phenyl, R⁶-heterocaryl, R⁶-naphthyl, C₃-C₁₀ cycloalkyl, (C₃-C₁₀)cycloalkyl/(C₁-C₆)alkyl or (C₁-C₆)alkoxy/(C₁-C₆)alkyl;

25 R²⁰, R²¹ and R²² are independently selected from the group consisting of H and C₁-C₆ alkyl; and

58

R²³ is C₁-C₆ alkyl or phenyl;

and wherein in the CCR5 antagonist compounds represented by the structural formula II:



or a pharmaceutically acceptable salt thereof, wherein

(1) X^e is -C(R¹³)₂-, -C(R¹³)(R¹⁹)-, -C(O)-, -O-, -NH-, -N((C₁-C₆)alkyl)-,

OR³ CH₂-(C₁-C₅)alkyl-R³ N(R⁴) O-(C₁-C₆)alkyl CH-(C₁-C₆)alkyl
-CR¹³-, -CR¹³-, -C-, -CR¹³-, -C-

10 O-C(O)-(C₁-C₆)alkyl O-C(O)-O-(C₁-C₆)alkyl O-C(O)-NH-(C₁-C₆)alkyl
-CR¹³-, -CR¹³-, -CR¹³-, -CR¹³-

O-C(O)-N((C₁-C₆)alkyl)₂ NR⁵-C(O)-(C₁-C₆)alkyl
-CR¹³-, -CR¹³-

15 NR⁵-C(O)-O-(C₁-C₆)alkyl NR⁵-C(O)-NH-(C₁-C₆)alkyl
-CR¹³-, -CR¹³-

NR⁵-C(O)-N((C₁-C₆)alkyl)₂ C(O)-(C₁-C₆)alkyl
-CR¹³-, or -N-


R⁶ is R^{6a}-phenyl, R^{6a}-pyridyl, R^{6a}-thiophenyl or R^{6b}-naphthyl;
R¹ is hydrogen, C₁-C₆ alkyl or C₂-C₆ alkenyl;

59

R² is R⁷, R⁸, R⁹-phenyl; R⁷, R⁸, R⁹-substituted 6-membered heteroaryl;

R⁷, R⁸, R⁹-substituted 6-membered heteroaryl N-oxide;

R¹⁰, R¹¹-substituted 5-membered heteroaryl; naphthyl; fluorenyl;



diphenylmethyl

5 R³ is R¹⁰,-phenyl, pyridyl, pyrroldyl, pyrazinyl or thiazolyl;

R⁴ is hydrogen, C₁-C₆ alkyl, fluoro-C₁-C₆ alkyl, cyclopropylmethyl,


-CH₂CH₂OH, -CH₂CH₂-O-(C₁-C₆)alkyl, -CH₂C(O)-O-(C₁-C₆)alkyl,

-CH₂C(O)NH₂, -CH₂C(O)-NH(C₁-C₆)alkyl or -CH₂C(O)-N(C₁-C₆)alkyl)₂;

R⁵ and R¹¹ are independently selected from the group consisting of

10 hydrogen and (C₁-C₆)-alkyl;

R^{6a} is 1 to 3 substituents independently selected from the group consisting of hydrogen, halogen, -CF₃, CF₃O-, -CN, -CF₃SO₂-, R¹²-phenyl,

-NHCOCF₃, 5-membered heteroaryl and , wherein X is -O-, -NH- or -N(CH₃)₂;

15 R⁶ is independently selected from the group consisting of R^{6a} and

CH₃SO₂-;

R⁷ and R⁸ are independently selected from the group consisting of (C₁-C₆)alkyl, halogen, -NR^{9a}R⁹¹, -OH, -CF₃, -OCH₃, -O-acyl, and -OCF₃;

R⁹ is R¹, hydrogen, phenyl, -NO₂, -CN, -CH₂F, -CHF₂, -CHO,

20 -CH=NOFR²⁰, pyridyl, pyridyl N-oxide, pyrrolidinyl, pyrazinyl,

-N(R²⁰)CONR²¹R²², -NHCONH(chloro-(C₁-C₆)alkyl), -NHCONH(C₃-C₁₀)-

cycloalkyl(C₁-C₆)alkyl), -NHCO(C₁-C₆)alkyl, -NHCOCF₃, -NHCO₂N(C₁-C₆)alkyl)₂,

NHSO₂(C₁-C₆)alkyl, -N(SO₂CF₃)₂, -NHSO₂(C₁-C₆)alkyl)₂,

-SO₂R²³, -SO₂CF₃, -SO₂NH(C₁-C₆ alkyl), -OSO₂(C₁-C₆)alkyl, -OSO₂CF₃, hydroxy(C₁-

25 C₆)alkyl, -CON R²⁰R²¹, -CON(CH₂CH₂-O-CH₃)₂,

-OCONH(C₁-C₆)alkyl, -CO₂R²⁰, -Si(CH₃)₃ or -B(OC(CH₃)₂)₂;

60

R¹⁰ is (C₁-C₆)alkyl, -NH₂ or R¹²-phenyl;

R¹² is 1 to 3 substituents independently selected from the group consisting

of hydrogen, (C₁-C₆) alkyl, -CF₃, -CO₂R²⁰, -CN, (C₁-C₆)alkoxy and halogen;

R¹³, R¹⁴, R¹⁵ and R¹⁶ are independently selected from the group

5 consisting of hydrogen and (C₁-C₆)alkyl;





R¹⁷ and R¹⁸ are independently selected from the group consisting of hydrogen and C₁-C₆ alkyl, or R¹⁷ and R¹⁸ together are a C₂-C₅ alkylene group and with the carbon to which they are attached form a spiro ring of 3 to 6 carbon atoms;

10 R⁹ is R²-phenyl, R⁹-heteroaryl, R⁹-naphthyl, C₃-C₁₀ cycloalkyl, (C³-C¹⁰)cycloalkyl(C₁-C₆)alkyl or (C₁-C₆)alkoxy(C₁-C₆)alkyl;


R²⁰, R²¹ and R²² are independently selected from the group consisting of H and C₁-C₆ alkyl; and

R²³ is C₁-C₆ alkyl or phenyl; or

15 (2):
Xe is -C(R¹³)(R¹⁹), -C(O)-, -O-, -NH-, -N(C₁-C₆)alkyl),



25



R^a is **R^{6b}**-phenyl, **R^{6b}**-pyridyl or **R^{6b}**-thiophenyl;


R4a is fluoro-C₁-C₆ alkyl, cyclopropylmethyl, -CH₂CH₂OH,

5 $\text{NH}-(\text{C}_1-\text{C}_6)_\text{alkyl}$ or $-\text{CH}_2\text{C}(\text{O})-\text{N}((\text{C}_1-\text{C}_6)_\text{alkyl})_2$;
 $-\text{CH}_2\text{CH}_2-\text{O}-(\text{C}_1-\text{C}_6)_\text{alkyl}$, $-\text{CH}_2\text{C}(\text{O})-\text{O}-(\text{C}_1-\text{C}_6)_\text{alkyl}$, $-\text{CH}_2\text{C}(\text{O})\text{NH}_2$, $-\text{CH}_2\text{C}(\text{O})-$

R^{8b} is CH₃SO₂-; and

$R^1, R^2, R^3, R^5, R^{14}, R^{15}, R^{16}$ and R^{19} are as defined in II(1).

and wherein in the CCR5 antagonist compounds represented by the structural formula III:




R is R⁸-phenyl, R⁸-pyridyl, R⁸-thiophenyl or R⁸-naphthyl;

R¹ is hydrogen or C₁-C₆ alkyl;

R² is R⁹, R¹⁰, R¹¹-phenyl; R⁹, R¹⁰, R¹¹-substituted 6-membered

15 heteroaryl; R⁹, R¹⁰, R¹¹-substituted 6-membered heteroaryl N-oxide;

R¹², R¹³-substituted 5-membered heteroaryl; naphthyl; fluorenyl;





R³ is hydrogen, C₁-C₆ alkyl, (C₁-C₆)alkoxy(C₁-C₆)alkyl, C₃-C₁₀ cycloalkyl, C₃-C₁₀ cycloalkyl/(C₁-C₆)alkyl, R⁸-phenyl, R⁸-phenyl/(C₁-C₆)alkyl, R⁸-naphthyl, R⁸ naphthyl/(C₁-C₆)alkyl, R⁸-heteroaryl or R⁸-heteroaryl/(C₁-C₆)alkyl;

R4, R5, R7 and R13 are independently selected from the group consisting of hydrogen and (C1-C6)-alkyl;


R⁶ is hydrogen, C₁-C₆ alkyl or C₂-C₆ alkenyl;

R⁸ is 1 to 3 substituents independently selected from the group consisting of hydrogen, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, -CF₃, CF₃O-, -CN, CH₃SO₂-, CF₃SO₂-, R¹⁴-phenyl, R¹⁴-benzyl, CH₃C(=NOCH₃),

5 $\text{CH}_3\text{C}(=\text{NOCH}_2\text{CH}_3)$, , $-\text{NH}_2$, $-\text{NHCOCF}_3$



-NHCONH(C₁-C₆ alkyl), -NHCO(C₁-C₆ alkyl), -NHSO₂(C₁-C₆ alkyl),

5-membered heterocaryl and , wherein X is -O-, -NH- or -N(CH₃)-;

R⁹ and R¹⁰ are independently selected from the group consisting of (C₁-C₆)alkyl, halogen, -NR¹¹R¹², -OH, -CF₃, -OCH₃, -O-acyl, -OCF₃ and

10 $-\text{Si}(\text{CH}_3)_3$

R¹¹ is R⁹, hydrogen, phenyl, -NO₂, -CN, -CH₂F, -CHF₂, -CHO, -CH=NOR¹⁷, pyridyl, pyridyl N-oxide, pyrimidinyl, pyrazinyl,

-N(R¹⁷)CONR¹⁸R¹⁹, -NHCONH(chloro-(C₁-C₈)alkyl), -NHCC

15 C_1 cycloalkyl/(C_1 - C_6)alkyl), -NHCO(C_1 - C_6)alkyl, -NHCOCF₃, -NH₂SO₂N((C_1 - C_6)alkyl)₂, -NH₂SO₂(C_1 - C_6)alkyl, -N(SO₂CF₃)₂, -NHCO₂(C_1 - C_6)alkyl, C_3 - C_{10} cycloalkyl, -SR²⁰, -SOR²⁰, -SO₂R²⁰, -SO₂NH(C_1 - C_6)alkyl), -OSO₂(C_1 - C_6)alkyl, -OSO₂CF₃, hydroxy(C_1 - C_6)alkyl, -CON R¹⁷, ¹⁸, -CON(CH₂CH₂-O-CH₃)₂, -OCONH(C_1 - C_6)alkyl, -CO₂R¹⁷, -Si(CH₃)₃ or -B(OC(CH₃)₂)₂;

R¹² is (C₁-C₆)alkyl, -NH₂ or R¹⁴-phenyl;

20 R¹⁴ is 1 to 3 substituents independently selected from the group consisting of hydrogen, (C₁-C₆) alkyl, -CF₃, -CO₂R¹⁷, -CN, (C₁-C₆)alkoxy and halogen;

R¹⁵ and R¹⁶ are independently selected from the group consisting of hydrogen and C₁-C₆ alkyl, or R¹⁵ and R¹⁶ together are a C₂-C₅ alkylene group and with the carbon to which they are attached form a spiro ring of 3 to 6 carbon atoms;

R^{17} , R^{18} and R^{19} are independently selected from the group consisting of H and C_1-C_6 alkyl; and


65

R¹⁵ and R¹⁶ are independently selected from the group consisting of hydrogen and C₁-C₆ alkyl; or R¹⁵ and R¹⁶ together are a C₂-C₅ alkylene group and with the carbon to which they are attached form a spiro ring of 3 to 6 carbon atoms;

5 R¹⁷, R¹⁸ and R¹⁹ are independently selected from the group consisting of H and C₁-C₆ alkyl; and

R²⁰ is C₁-C₆ alkyl or phenyl; or

(2) R^a is R⁸-phenyl, R⁸-pyridyl or R⁸-thiophenyl;



10 and R¹, R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸, R¹⁹ and R²⁰ are as defined in IV(1).

3 The use of any preceding claim, wherein the patients are treatment-naïve or treatment-experienced patients.

15

4 The use of any preceding claim, wherein the patients are treatment-naïve or treatment-experienced pediatric patients.

5 The use of any preceding claim, wherein the pegylated interferon- α a administered is pegylated interferon α a-2a or pegylated interferon α a-2b.

20

6 The use of any preceding claim, wherein the treatment comprises a therapeutically effective amount of pegylated interferon- α a in association with a therapeutically effective amount of a CCR5 antagonist sufficient to lower HIV-1-RNA levels in such patients.

25

7 The use of any preceding claim, wherein the pegylated interferon- α a administered is a pegylated interferon α a-2b and wherein the amount of

66

pegylated interferon α a-2b administered is in the range of about 0.1 to about 9.0 micrograms per kilogram per week, preferably in the range of about 0.5 to about 3.0 micrograms per kilogram per week, and more preferably is in the range of about 0.75 to about 1.5 micrograms per kilogram per week

5

8 The use of any preceding claim, wherein the pegylated interferon- α a administered is a pegylated interferon α a-2a and the amount of pegylated interferon α a-2a administered is in the range of about 50 to about 500 micrograms per week, preferably in the range of about 150 to about 250 micrograms per week, or preferably in the range of about 180 to about 250 micrograms per week, and more preferably is in the range of about 150 to about 180 micrograms once per week.

10

9 The use of any preceding claim, wherein the treatment further comprises a therapeutically effective amount of at least one of ribavirin, IL-2, IL-12, pentastatin alone or in combination with a therapeutically effective amount of an anti-HIV-1 drug therapy.

15



10 The use of any preceding claim wherein the patients are co-infected with HIV-1 and HCV and wherein the treatment further comprises a therapeutically effective amount of ribavirin and a therapeutically effective amount of an anti-HIV-1 drug therapy.

20

11 The use of claim 9 or 10 wherein the anti-HIV-1 drug therapy is HAART.


25

12 The use of a pegylated interferon- α a, a CCR5 antagonist, ribavirin and HAART for the preparation of a medicament for the treatment of HIV-1 and HCV co-infections in patients wherein the CCR5 antagonist is represented by the structural formula I or II or III or IV:




5 or a pharmaceutically acceptable salt of I or II or III or IV; sufficient to lower HIV-1-RNA and HCV-RNA levels;

wherein in the CCR5 antagonist compounds represented by structural formula 1:




X is -C(R¹³)₂-, -C(R¹³)(R¹⁹)-, -C(O)-, -O-, -NH-, -N((C₁-C₆alkyl))-,



WO 00/66141

PCT/US00/11634




R is R⁶-phenyl, R⁶-pyridyl, R⁶-thiophenyl or R⁶-naphthyl;

R¹ is hydrogen, C₁-C₆ alkyl or C₂-C₆ alkenyl;

R² is R⁷, R⁸, R⁹-phenyl; R⁷, R⁸, R⁹-substituted 6-membered heterocaryl;

R⁷, R⁸, R⁹-substituted 6-membered heteroaryl N-oxide;

10 R¹⁰, R¹¹-substituted 5-membered heteroaryl; naphthyl; fluorenyl;



R³ is R⁶-phenyl, R⁶-heteroaryl or R⁶-naphthyl;

R⁴ is hydrogen, C₁-C₆ alkyl, fluoro-C₁-C₆ alkyl, cyclopropylmethyl,


$$-\text{CH}_2\text{CH}_2\text{OH}, -\text{CH}_2\text{CH}_2-\text{O}-(\text{C}_1-\text{C}_6)\text{alkyl}, -\text{CH}_2\text{C}(\text{O})-\text{O}-(\text{C}_1-\text{C}_6)\text{alkyl},$$

15 $-\text{CH}_2\text{C}(\text{O})\text{NH}_2$, $-\text{CH}_2\text{C}(\text{O})-\text{NH}(\text{C}_1\text{-C}_6\text{alkyl})$ or $-\text{CH}_2\text{C}(\text{O})-\text{N}(\text{C}_1\text{-C}_6\text{alkyl})_2$.

R⁵ and R¹¹ are independently selected from the group consisting of hydrogen and (C₁-C₆)-alkyl;

R⁶ is 1 to 3 substituents independently selected from the group consisting of hydrogen, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, -CF₃, CF₃O-, CH₂C(O)-, -CN,

20 CH_3SO_2 -, CF_3SO_2 -, R^{14} -phenyl, R^{14} -benzyl, $\text{CH}_3\text{C}(=\text{NOCH}_3)$ -,



69

NHCO(C₁-C₆ alkyl), -NHSO₂(C₁-C₆ alkyl),



5-membered heterocycle and , wherein X is -O-, -NH- or -N(CH₃)-;

R⁷ and R⁸ are independently selected from the group consisting of (C₁-C₆)alkyl, halogen, -NR²⁰R²¹, -OH, -CF₃, -OCH₃, -O-acyl, and -OCF₃;

5 (C₆)alkyl, halogen, -NR²⁰R²¹, -OH, -CF₃, -OCH₃, -O-acyl, and -OCF₃;

R⁸ is R⁷, hydrogen, phenyl, -NO₂, -CN, -CH₂F, -CHF₂, -CHO, -CH=NO²⁰, pyridyl, pyridyl N-oxide, pyrimidinyl, pyrazinyl,

-N(R²⁰)CONR²¹R²², -NHCONH(chloro-(C₁-C₆)alkyl), -NHCONH((C₃-C₁₀)-cycloalkyl)(C₁-C₆)alkyl), -NHCO(C₁-C₆)alkyl, -NHCOCF₃, -NHSO₂N((C₁-C₆)alkyl)₂, -

NHSO₂(C₁-C₆)alkyl, -N(SO₂CF₃)₂, -NHCO₂(C₁-C₆)alkyl, C₃-C₁₀ cycloalkyl, -SR²³, -

SOR²³, -SO₂R²³, -SO₂NH(C₁-C₆ alkyl), -OSO₂(C₁-C₆)alkyl, -OSO₂CF₃, hydroxy(C₁-C₆)alkyl, -CON R²⁰R²¹, -CON(CH₂CH₂O-CH₃)₂,

-OCONH(C₁-C₆)alkyl, -CO₂R²⁰, -Si(CH₃)₃ or -B(OC(CH₃)₂)₂;

10 R¹⁰ is (C₁-C₆)alkyl, -NH₂ or R¹²-phenyl;

R¹² is 1 to 3 substituents independently selected from the group consisting of hydrogen, (C₁-C₆) alkyl, -CF₃, -CO₂R₂₀, -CN, (C₁-C₆)alkoxy and halogen;

15 R¹³, R¹⁴, R¹⁵ and R¹⁶ are independently selected from the group consisting of hydrogen and (C₁-C₆)alkyl;

R¹⁷ and R¹⁸ are independently selected from the group consisting of hydrogen and C₁-C₆ alkyl, or R¹⁷ and R¹⁸ together are a C₂-C₅ alkylene group and with the carbon to which they are attached form a spiro ring of 3 to 6 carbon atoms;

20 R¹⁹ is R⁶-phenyl, R⁶-heteroaryl, R⁶-naphthyl, C₃-C₁₀ cycloalkyl, (C₃-C₁₀)cycloalkyl(C₁-C₆)alkyl or (C₁-C₆)alkoxy(C₁-C₆)alkyl;

R²⁰, R²¹ and R²² are independently selected from the group consisting of H and C₁-C₆ alkyl; and

R²³ is C₁-C₆ alkyl or phenyl;

and wherein in the CCR5 antagonist compounds represented by the structural formula II:

structural formula II:

structural formula II:

structural formula II:

structural formula II:

structural formula II:

structural formula II:

structural formula II:


structural formula II:

structural formula II:

structural formula II:

structural formula II:

70




71

R⁴ is hydrogen, C₁-C₆ alkyl, fluoro-C₁-C₆ alkyl, cyclopropylmethyl, -CH₂CH₂OH, -CH₂CH₂-O-(C₁-C₆)alkyl, -CH₂C(O)-O-(C₁-C₆)alkyl, -CH₂C(O)NH₂, -CH₂C(O)-NH(C₁-C₆)alkyl or -CH₂C(O)-N((C₁-C₆)alkyl)₂;

R⁵ and R¹¹ are independently selected from the group consisting of hydrogen and (C₁-C₆)-alkyl;

R^{6a} is 1 to 3 substituents independently selected from the group consisting of hydrogen, halogen, -CF₃, CF₃O-, -CN, -CF₃SO₂, R¹²-phenyl,

-NHCOCF₃, 5-membered heteroaryl and , wherein X is -O-, -NH- or -N(CH₃)-;

10. R⁶ is independently selected from the group consisting of R^{6a} and CH₃SO₂-;

R⁷ and R⁸ are independently selected from the group consisting of (C₁-C₆)alkyl, halogen, -NR^{9a}R¹¹, -OH, -CF₃, -OCH₃, -O-acyl, and -OCF₃;

15. R⁹ is R¹, hydrogen, phenyl, -NO₂, -CN, -CH₂F, -CHF₂, -CHO, -CH=NOR²⁰, pyridyl, pyridyl N-oxide, pyrimidinyl, pyrazinyl, -N(R²⁰)CONR²¹R²², -NHCONH(chloro-(C₁-C₆)alkyl), -NHCONH((C₃-C₁₀)-cycloalkyl((C₁-C₆)alkyl), -NHCO(C₁-C₆)alkyl, -NHCOCF₃, -NHCO₂N((C₁-C₆)alkyl)₂, -NHCO₂(C₁-C₆)alkyl, -N(SO₂CF₃)₂, -NHCO₂(C₁-C₆)alkyl, C₃-C₁₀ cycloalkyl, -SR²³, -SOR²³, -SO₂R²³, -SO₂NH(C₁-C₆ alkyl), -OSO₂(C₁-C₆)alkyl, -OSO₂CF₃, hydroxy(C₁-C₆)alkyl, -CON R²⁹R²¹, -CON(CH₂CH₂-O-CH₃)₂, -OCONH(C₁-C₆)alkyl, -CO₂R²⁰, -Si(CH₃)₂ or -B(OC(CH₃)₂)₂;

R¹⁰ is (C₁-C₆)alkyl, -NH₂ or R¹²-phenyl;

R¹² is 1 to 3 substituents independently selected from the group consisting of hydrogen, (C₁-C₆) alkyl, -CF₃, -CO₂R²⁰, -CN, (C₁-C₆)alkoxy and halogen;

25. R¹³, R¹⁴, R¹⁵ and R¹⁶ are independently selected from the group consisting of hydrogen and (C₁-C₆)alkyl;

R¹⁷ and R¹⁸ are independently selected from the group consisting of hydrogen and C₁-C₆ alkyl, or R¹⁷ and R¹⁸ together are a C₂-C₅ alkylene group

72

and with the carbon to which they are attached form a spiro ring of 3 to 6 carbon atoms;


R¹⁹ is R⁶-phenyl, R⁶-heteroaryl, R⁶-naphthyl, C₃-C₁₀ cycloalkyl, (C₃-C₁₀)cycloalkyl((C₁-C₆)alkyl) or (C₁-C₆)alkoxy(C₁-C₆)alkyl;


5. R²⁰, R²¹ and R²² are independently selected from the group consisting of H and C₁-C₆ alkyl; and


R²³ is C₁-C₆ alkyl or phenyl; or


(2):

X^e is -C(R¹³)(R¹⁹), -C(O)-, -O-, -NH-, -N((C₁-C₆)alkyl)-,

10. 



15. 





20. R^e is R^{6b}-phenyl, R^{6b}-pyridyl or R^{6b}-thiophenyl;

R^{4e} is fluoro-C₁-C₆ alkyl, cyclopropylmethyl, -CH₂CH₂OH,


-CH₂CH₂-O-(C₁-C₆)alkyl, -CH₂C(O)-O-(C₁-C₆)alkyl, -CH₂C(O)NH₂, -CH₂C(O)-NH-(C₁-C₆)alkyl or -CH₂C(O)-N((C₁-C₆)alkyl)₂;

R^{6b} is CH₃SO₂-; and

73

R¹, R², R³, R⁵, R¹⁴, R¹⁵, R¹⁶ and R¹⁸ are as defined in II(1);

and wherein in the CCR5 antagonist compounds represented by the structural formula III:



5


R is R⁸-phenyl, R⁸-pyridyl, R⁸-thiophenyl or R⁸-naphthyl;

R¹ is hydrogen or C₁-C₆ alkyl;

R² is R⁹, R¹⁰, R¹¹-phenyl; R⁹, R¹⁰, R¹¹-substituted 6-membered

heteroaryl; R⁹, R¹⁰, R¹¹-substituted 6-membered heteroaryl N-oxide;

10 R¹², R¹³-substituted 5-membered heteroaryl; naphthyl; fluorenyl;



R³ is hydrogen, C₁-C₆ alkyl, (C₁-C₆)alkoxy(C₁-C₆)alkyl, C₃-C₁₀ cycloalkyl,


C₃-C₁₀ cycloalkyl(C₁-C₆)alkyl, R⁸-phenyl, R⁸-phenyl(C₁-C₆)alkyl, R⁸-naphthyl, R⁸-naphthyl(C₁-C₆)alkyl, R⁸-heteroaryl or R⁸-heteroaryl(C₁-C₆)alkyl;

15 R⁴, R⁵, R⁷ and R¹³ are independently selected from the group consisting of hydrogen and (C₁-C₆)-alkyl;

R⁶ is hydrogen, C₁-C₆ alkyl or C₂-C₆ alkenyl;

R⁸ is 1 to 3 substituents independently selected from the group consisting of hydrogen, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, -CF₃, CF₃O-, CH₃C(O)-, -CN,

20 CH₃SO₂-, CF₃SO₂-, R¹⁴-phenyl, R¹⁴-benzyl, CH₃C(=NOCH₃),



74

-NHCONH(C₁-C₆ alkyl), -NHCO(C₁-C₆ alkyl), -NHSO₂(C₁-C₆ alkyl),

5-membered heteroaryl and , wherein X is -O-, -NH- or -N(CH₃)-;

R⁹ and R¹⁰ are independently selected from the group consisting of (C₁-C₆)alkyl, halogen, -NR¹⁷R¹⁸, -OH, -CF₃, -OCH₃, -O-acyl, -OCF₃ and

5 -Si(CH₃)₃;

R¹¹ is R⁹, hydrogen, phenyl, -NO₂, -CN, -CH₂F, -CHF₂, -CHO,

-CH=NOH¹⁷, pyridyl, pyridyl N-oxide, pyrimidinyl, pyrazinyl,

-N(R¹⁷)CONH(R¹⁸)R¹⁹, -NHCONH(chloro-(C₁-C₆)alkyl), -NHCONH((C₃-

C₁₀)cycloalkyl(C₁-C₆)alkyl), -NHCO(C₁-C₆)alkyl, -NHCOCF₃, -NHSO₂N((C₁-

10 C₆)alkyl)₂, -NHSO₂(C₁-C₆)alkyl, -N(SO₂CF₃)₂, -NHCO₂(C₁-C₆)alkyl, C₃-C₁₀

cycloalkyl, -SR²⁰, -SOR²⁰, -SO₂R²⁰, -SO₂NH(C₁-C₆ alkyl), -OSO₂(C₁-C₆)alkyl, -

OSO₂CF₃, hydroxy(C₁-C₆)alkyl, -CON R¹⁷R¹⁸, -CON(CH₂CH₂O-CH₃)₂, -

OCONH(C₁-C₆)alkyl, -CO₂R¹⁷, -Si(CH₃)₃ or -B(OC(CH₃)₂)₂;

R¹² is (C₁-C₆)alkyl, -NH₂ or R¹⁴-phenyl;

15 R¹⁴ is 1 to 3 substituents independently selected from the group consisting of hydrogen, (C₁-C₆) alkyl, -CF₃, -CO₂R¹⁷, -CN, (C₁-C₆)alkoxy and halogen;

R¹⁵ and R¹⁶ are independently selected from the group consisting of

hydrogen and C₁-C₆ alkyl, or R¹⁵ and R¹⁶ together are a C₂-C₆ alkylene group and with the carbon to which they are attached form a spiro ring of 3 to 6 carbon

20 atoms;


R¹⁷, R¹⁸ and R¹⁹ are independently selected from the group consisting of H

and C₁-C₆ alkyl; and

R²⁰ is C₁-C₆ alkyl or phenyl;

25 and wherein in the CCR5 antagonist compounds represented by the structural formula IV:

75



or a pharmaceutically acceptable salt thereof, wherein


(1) R^a is R^{8a} -phenyl, R^{8b} -pyridyl, R^{8b} -thiophenyl or R^8 -naphthyl;

R^1 is hydrogen or C_1 - C_6 alkyl;

5 R^2 is R^9 , R^{10} , R^{11} -phenyl; R^9 , R^{10} , R^{11} -substituted 6-membered

heteroaryl; R^9 , R^{10} , R^{11} -substituted 6-membered heteroaryl N-oxide;

R^{12} , R^{13} -substituted 5-membered heteroaryl; naphthyl; fluorenyl;




R^3 is hydrogen, C_1 - C_6 alkyl, (C_1-C_6) alkoxy (C_1-C_6) alkyl, C_3-C_{10} cycloalkyl,

10 C_3-C_{10} cycloalkyl (C_1-C_6) alkyl, R^8 -phenyl, R^8 -phenyl (C_1-C_6) alkyl, R^8 -naphthyl, R^8 -naphthyl (C_1-C_6) alkyl, R^8 -heteroaryl or R^8 -heteroaryl (C_1-C_6) alkyl;

R^4 , R^5 , R^7 and R^{13} are independently selected from the group consisting of hydrogen and (C_1-C_6) -alkyl;

R^6 is hydrogen, C_1 - C_6 alkyl or C_2 - C_6 alkenyl;

15 R^8 is 1 to 3 substituents independently selected from the group consisting of hydrogen, halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-CF_3$, CF_3O , $CH_3C(O)$ -, $-CN$, CH_3SO_2 -, CF_3SO_2 -, R^{14} -phenyl, R^{14} -benzyl, $CH_3C(=NOCH_3)$,




$CH_3C(=NOCH_2CH_3)$, $-NH_2$, $-NHCOCF_3$,

20 $-NHCONH(C_1-C_6)$ alkyl), $-NHCO(C_1-C_6)$ alkyl), $-NHCO_2(C_1-C_6)$ alkyl), $-N(CH_3)$;



76


R^{8a} is 1 to 3 substituents independently selected from the group consisting of hydrogen, halogen, $-CF_3$, CF_3O -, $-CN$, CF_3SO_2 -, R^{14} -phenyl, $-NHCOCF_3$, 5-


membered heteroaryl and , wherein X is as defined above;

R^{8b} is 1 to 3 substituents independently selected from the group consisting

5 of hydrogen, halogen, $-CF_3$, CF_3O -, $CH_3C(O)$ -, $-CN$, CF_3SO_2 -, R^{14} -benzyl,

$CH_3C(=NOCH_3)$, $CH_3C(=NOCH_2CH_3)$,



$-NHCOCF_3$, 5-membered heteroaryl and , wherein X is as defined above;

R^9 and R^{10} are independently selected from the group consisting of $(C_1$ -

$C_6)$ alkyl, halogen, $-NR^{17}R^{18}$ -, $-OH$ -, $-CF_3$ -, $-OCH_3$ -, $-O$ -acyl-, $-OCF_3$ and

10 $-Si(CH_3)_3$;

R^{11} is R^9 , hydrogen, phenyl, $-NO_2$ -, $-CN$ -, $-CH_2F$ -, $-CHF_2$ -, $-CHO$,

$-CH=NOH$ -, $-CH=NOH$ -, $-CH=NOH$ -, $-CH=NOH$ -, $-CH=NOH$ -, $-CH=NOH$ -, $-CH=NOH$ -, $-CH=NOH$ -,

$-N(R^{17})CONR^{18}R^{19}$ -, $-NHCONH(chloro-(C_1-C_6)$ alkyl), $-NHCONH((C_3$ -

$C_1)$ cycloalkyl (C_1-C_6) alkyl), $-NHCO(C_1-C_6)$ alkyl, $-NHCOCF_3$ -, $-NHCO_2N((C_1$ -

15 $C_6)$ alkyl) $_2$ -, $-NHCO_2(C_1-C_6)$ alkyl, $-N(SO_2CF_3)_2$ -, $-NHCO_2(C_1-C_6)$ alkyl, C_3-C_{10}

cycloalkyl, $-SR^{20}$ -, $-SOR^{20}$ -, $-SO_2R^{20}$ -, $-SO_2NH(C_1-C_6)$ alkyl), $-OSO_2(C_1-C_6)$ alkyl, $_2$ -,

$-OSO_2CF_3$ -, hydroxy (C_1-C_6) alkyl, $-CONR^{17}R^{18}$ -, $-CON(CH_2CH_2O-CH_3)_2$ -,

$-OCONH(C_1-C_6)$ alkyl, $-CO_2R^{17}$ -, $-Si(CH_3)_3$ or $-B(OC(CH_3)_2)_2$;

R^{12} is (C_1-C_6) alkyl, $-NH_2$ or R^{14} -phenyl;

20 R^{14} is 1 to 3 substituents independently selected from the group consisting of hydrogen, (C_1-C_6) alkyl, $-CF_3$ -, $-CO_2R^{17}$ -, $-CN$ -, (C_1-C_6) alkoxy and halogen;

R^{15} and R^{16} are independently selected from the group consisting of

hydrogen and C_1 - C_6 alkyl, or R^{15} and R^{16} together are a C_2 - C_5 alkylene group

and with the carbon to which they are attached form a spiro ring of 3 to 6 carbon

25 atoms;

77

R^{17} , R^{18} and R^{19} are independently selected from the group consisting of H and C_1-C_6 alkyl; and

R^{20} is C_1-C_6 alkyl or phenyl; or

(2) R^a is R^8 -phenyl, R^8 -pyridyl or R^8 -thiophenyl;



and R^1 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} and R^{20} are as defined in IV(1).

10 13. The use of claim 12, wherein the patients are treatment- experienced patients or treatment-naïve patients.

14. The use of claim 12, wherein the patients are treatment- experienced patients or treatment-naïve pediatric patients.

15 15. The use of claim 12, wherein the pegylated interferon- α a administered is pegylated interferon α a-2a or pegylated interferon α a-2b.

16. The use of claim 15, wherein the pegylated interferon- α a administered is a pegylated interferon α a-2b and wherein the amount of pegylated interferon α a-2b administered is in the range of about 0.1 to about 9.0 micrograms per kilogram per week, preferably in the range of about 0.5 to about 3.0 micrograms per kilogram per week, and more preferably is in the range of about 0.75 to about 1.5 micrograms per kilogram per week.

25 17. The use of claim 15, wherein the pegylated interferon- α a administered is a pegylated interferon α a-2a and the amount of pegylated interferon α a-2a administered is in the range of about 50 to about 500 micrograms per week, preferably in the range of about 150 to about

78

250 micrograms per week, and more preferably in the range of about 180 to about 250 micrograms per week, or preferably is in the range of about 150 to about 180 micrograms per week.

5 18. The use of claim 12, wherein the patient is a pediatric patient and the amount of ribavirin is in a range of about 8 to about 15 mg per kilogram per day, in divided doses.

10 19. The use of claim 12, wherein the amount of ribavirin is from about 400 to about 1600 mg per day, preferably about 600 to about 1200 mg/day or about 800 to about 1200 mg day and most preferably about 1000 to about 1200 mg/kg a day.

15

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 November 2000 (09.11.2000)

PCT

(10) International Publication Number
WO 00/66141 A3

(31) International Patent Classification: A61K 38/21, A61P 31/18 // (A61K 38/21, 31:4545) (A61K 38/21, 31:496)

(74) Agents: HOFFMANN, Thomas, D. et al.; Schering-Plough Corporation, Patent Department, K-6-1 1990, 2000 Gallop-Hill Road, Kenilworth, NJ 07033-0530 (US).

(21) International Application Number: PCT/US00/1634

(22) International Filing Date: 1 May 2000 (01.05.2000)

(23) Filing Language: English

(26) Publication Language: English

(30) Priority Date: 4 May 1999 (04.05.1999) US 09/304,897

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application: 09/304,897 (CIP) US Filed on 4 May 1999 (04.05.1999)

(71) Applicant (for all designated States except US): SCHERING CORPORATION (US/US); 2000 Gallop-Hill Road, Kenilworth, NJ 07033-0530 (US).

(72) Inventor and Inventor/Applicant (for US only): LAUGHLIN, Mark, A. (US/US); 25 Cinder Road, #3M, Edison, NJ 08820 (US).

(88) Date of publication of the international search report: 8 February 2001

(84) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GR, HU, ID, IL, IN, JP, KG, KP, KZ, LC, LK, LR, LT, LU, LV, MA, MD, MG, MK, MN, MX, NO, NZ, PL, PT, RO, RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, US, UZ, VN, YU, ZA.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

INTERNATIONAL SEARCH REPORT

Internat. Application No
PCT/US 00/1634

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K38/21 A61P31/18 // (A61K38/21, 31:4545), (A61K38/21, 31:496)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELD OF SEARCHED
Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data bases consulted during the international search (name of data base and, where practical, search terms used)
EPO-Internal, WPI Data, BIOSIS, MEDLINE, EMBASE, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 25617 A (SPRINGER MARTIN S.; MACCOSS MALCOLM (US); MERCK & CO INC (US); MILL) 18 June 1998 (1998-06-18) abstract page 5, line 6 - line 14	1,3-11
Y	WO 99 17773 A (PORTER RODERICK A.; SMITHLINE BEECHAM PLC (GB); BONDINELL WILLIAM) 15 April 1999 (1999-04-15) the whole document	1,3-11
Y	WO 98 48840 A (SCHERING CORP) 5 November 1998 (1998-11-05) the whole document	1,3-11
A	WO 97 42969 A (HOFFMANN LA ROCHE) 20 November 1997 (1997-11-20) the whole document	

* Further documents are listed in the continuation of box C. ☒ Patent family members are listed in annex.

* Special categories of cited documents:

- * document defining the general state of the art which is not considered to be of particular relevance
- * earlier document but published on or after the international filing date which may, however, be of particular relevance
- * document which may, however, be of particular relevance but which is cited to establish the publication date of another document or other special reason (as specified)
- * document relating to an oral disclosure, use, exhibition or other means
- * document published prior to the international filing date but later than the priority date claimed
- * document member of the same patent family

Date of the actual completion of the international search

27 November 2000

05/12/2000

Name and mailing address of the ISA
European Patent Office, P.O. Box 5518, Postfach 2
NL - 2200 HV Rijswijk
Tel. (+31-70) 340-2000, Tx. 31 651 apo nl,
Fax (+31-70) 340-3010

Authorized officer
Niemann, F

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No.
PCT/US 00/11634

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9825617 A	18-06-1998	AU 5522498 A	03-07-1998
WO 9917773 A	15-04-1999	AU 9790198 A	27-04-1999
		EP 1037635 A	27-09-2000
		ZA 9809083 A	07-04-1999
WO 9848840 A	05-11-1998	US 5908621 A	01-06-1999
		AU 7249098 A	24-11-1998
		BR 9809425 A	25-07-2000
		CN 1261806 T	02-08-2000
		EP 0975369 A	02-02-2000
		NO 995263 A	27-12-1999
		PL 336576 A	03-07-2000
		ZA 9803543 A	10-11-1998
WO 9742969 A	20-11-1997	BR 9709238 A	10-08-1999
		CN 1218410 A	02-06-1999
		EP 0907373 A	14-04-1999
		JP 2000504029 T	04-04-2000
		US 5928636 A	27-07-1999

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☒ OTHER: side text

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.